

# INTERSECTIONS OF SCHUBERT VARIETIES AND EIGENVALUE INEQUALITIES IN AN ARBITRARY FINITE FACTOR

H. BERCOVICI, B. COLLINS, K. DYKEMA, W. S. LI, AND D. TIMOTIN

**ABSTRACT.** It is known that the eigenvalues of selfadjoint elements  $a, b, c$  with  $a + b + c = 0$  in the factor  $\mathcal{R}^\omega$  are characterized by a system of inequalities analogous to the classical Horn inequalities of linear algebra. We prove that these inequalities are in fact true for elements of an arbitrary finite factor. In particular, if  $x, y, z$  are selfadjoint elements of such a factor and  $x + y + z = 0$ , then there exist selfadjoint  $a, b, c \in \mathcal{R}^\omega$  such that  $a + b + c = 0$  and  $a$  (respectively,  $b, c$ ) has the same eigenvalues as  $x$  (respectively,  $y, z$ ). A ('complete') matricial form of this result is known to imply an affirmative answer to an embedding question formulated by Connes.

The main difficulty in our argument is the proof that certain generalized Schubert cells (consisting of projections of fixed trace) have nonempty intersection. In finite dimensions, this follows from classical intersection theory. Our approach is to exhibit an actual element in this intersection, given by a formula which applies generically regardless of the algebra (or of the dimension of the underlying space). This argument requires a good understanding of the combinatorial structure of honeycombs, and it seems to be new even in finite dimensions.

## INTRODUCTION

Assume that  $A, B, C$  are complex selfadjoint  $n \times n$  matrices, and  $A + B + C = 0$ . A. Horn proposed in [19] the question of characterizing the possible eigenvalues of these matrices, and indeed he conjectured an answer which was eventually proved correct due to efforts of A. Klyachko [20] and A. Knutson and T. Tao [21]. To explain this characterization, list the eigenvalues of  $A$ , repeated according to multiplicity, in nonincreasing order

$$\lambda_A(1) \geq \lambda_A(2) \geq \cdots \geq \lambda_A(n),$$

choose an orthonormal basis  $x_j \in \mathbb{C}^n$  such that  $Ax_j = \lambda_A(j)x_j$ , and denote by  $E_A(j)$  the space generated by  $\{x_1, x_2, \dots, x_j\}$ . Horn's conjecture involves, in addition to the trace identity

$$\sum_{j=1}^n (\lambda_A(j) + \lambda_B(j) + \lambda_C(j)) = 0,$$

---

1991 *Mathematics Subject Classification.* Primary: 14N15; Secondary: 15A42, 46L10, 46L54, 52B05, 05E99.

HB, KD, and WSL were supported in part by grants from the National Science Foundation. BC was supported in part by an NSERC grant.

a collection of inequalities of the form

$$\sum_{i \in I} \lambda_A(i) + \sum_{j \in J} \lambda_B(j) + \sum_{k \in K} \lambda_C(k) \leq 0,$$

where  $I, J, K \subset \{1, 2, \dots, n\}$  are sets with equal cardinalities. One way to prove such inequalities is to observe that

$$\mathrm{Tr}(PAP + PBP + PCP) = 0$$

for any orthogonal projection  $P$ , and to find a projection  $P$  such that

$$\mathrm{Tr}(PAP) \geq \sum_{i \in I} \lambda_A(i), \quad \mathrm{Tr}(PBP) \geq \sum_{j \in J} \lambda_B(j), \quad \mathrm{Tr}(PCP) \geq \sum_{k \in K} \lambda_C(k).$$

Now, if  $I = \{i_1 < i_2 < \dots < i_r\}$ , the first condition is guaranteed provided that the range  $M$  of  $P$  has dimension  $r$ , and

$$\dim(M \cap E_A(i_\ell)) \geq \ell, \quad \ell = 1, 2, \dots, r.$$

These conditions describe the Schubert variety  $S(E_A, I)$  determined by the flag  $\{E_A(\ell)\}_{\ell=1}^n$  and the set  $I$ . Thus, such a projection can be found provided that

$$S(E_A, I) \cap S(E_B, J) \cap S(E_C, K) \neq \emptyset.$$

Klyachko [20] proved that the collection of all inequalities obtained this way is sufficient to answer Horn's question, and observed that Horn's conjectured answer would also be proved if a certain 'saturation conjecture' were true. This conjecture was proved by Knutson and Tao [21]. (See also [22] for a direct proof of Horn's conjecture, and [15] for a very good survey of the history of the problem and its ramifications. Some earlier expositions are in [11, 14].)

There are several infinite-dimensional analogues of the Horn problem. One can for instance consider compact selfadjoint operators  $A, B, C$  on a Hilbert space and their eigenvalues. This analogue was considered by several authors [13, 16], and a complete solution can be found in [5] for operators such that  $A, B$ , and  $-C$  are positive, and [6] for the general case. Without going into detail, let us say that these solutions are based on an understanding of the behavior of the Horn inequalities as the dimension of the space tends to infinity. The analogue we are interested in here replaces the algebra  $M_n(\mathbb{C})$  of  $n \times n$  matrices by a finite factor. This is simply a selfadjoint algebra  $\mathcal{A}$  of operators on a complex Hilbert space  $H$  such that  $\mathcal{A}' \cap \mathcal{A} = \mathbb{C}1_H$  (where  $\mathcal{A}' = \{T : AT = TA \text{ for all } A \in \mathcal{A}\}$ ),  $\mathcal{A}'' = \mathcal{A}$ , and for which there exists a linear functional  $\tau : \mathcal{A} \rightarrow \mathbb{C}$  such that  $\tau(X^*X) = \tau(XX^*) > 0$  for all  $X \in \mathcal{A} \setminus \{0\}$ . The algebras  $M_n(\mathbb{C})$  are finite factors. When  $\mathcal{A}$  is an infinite dimensional finite factor, it is called a factor of type  $\mathrm{II}_1$ . A complete flag in a  $\mathrm{II}_1$  factor  $\mathcal{A}$  is a family of orthogonal projections  $\{E(t) : 0 \leq t \leq \tau(1_H)\}$  such that  $\tau(E(t)) = t$ , and  $E(t) \leq E(s)$  for  $t \leq s$ . For any selfadjoint operator  $A \in \mathcal{A}$  there exist a nonincreasing function  $\lambda_A : [0, \tau(1_H)] \rightarrow \mathbb{R}$ , and a complete flag  $\{E_A(t) : 0 \leq t \leq \tau(1_H)\}$  such that

$$A = \int_0^{\tau(1_H)} \lambda_A(t) dE_A(t).$$

This is basically a restatement of the spectral theorem. The function  $\lambda_A$  is uniquely determined at its points of continuity, but the space  $E_A(t)$  is not uniquely determined on the open intervals where  $\lambda_A$  is constant. Note that

$$\tau(A) = \int_0^{\tau(1_H)} \lambda_A(t) dt,$$

and therefore we have a trace identity

$$\int_0^{\tau(1_H)} (\lambda_A(t) + \lambda_B(t) + \lambda_C(t)) dt = 0$$

whenever  $A + B + C = 0$ .

Many factors of type  $\text{II}_1$  can be approximated in a weak sense by matrix algebras. These are the factors that embed in the ultrapower  $\mathcal{R}^\omega$  of the hyperfinite  $\text{II}_1$  factor  $\mathcal{R}$ . For elements in such factors one can easily prove analogues of Horn's inequalities. More precisely, assume that  $\lambda : [0, T] \rightarrow \mathbb{R}$  is a nonincreasing function. The sequence

$$\lambda^{(n)}(1) \geq \lambda^{(n)}(2) \geq \dots \geq \lambda^{(n)}(n)$$

is defined by  $\lambda^{(n)}(j) = \int_{(j-1)T/n}^{jT/n} \lambda(t) dt$ . The following result was proved in [4]. We use the normalization  $\tau(1_H) = 1$  for the factor  $\mathcal{R}^\omega$ .

**Theorem 0.1.** *Let  $\alpha, \beta, \gamma : [0, 1] \rightarrow \mathbb{R}$  be nonincreasing functions. The following are equivalent:*

- (1) *There exist selfadjoint operators  $A, B, C \in \mathcal{R}^\omega$  such that  $\lambda_A = \alpha, \lambda_B = \beta, \lambda_C = \gamma$ , and  $A + B + C = 0$ .*
- (2) *For every integer  $n \geq 1$ , there exist matrices  $A_n, B_n, C_n \in M_n(\mathbb{C})$  such that  $\lambda_{A_n} = \alpha^{(n)}, \lambda_{B_n} = \beta^{(n)}, \lambda_{C_n} = \gamma^{(n)}$ , and  $A_n + B_n + C_n = 0$ .*

Note that condition (2) above requires, in addition to the trace identity, an infinite (and infinitely redundant) collection of Horn inequalities. We will show that these inequalities are in fact satisfied in any factor of type  $\text{II}_1$ .

**Theorem 0.2.** *Given a factor  $\mathcal{A}$  of type  $\text{II}_1$ , selfadjoint elements  $A, B, C \in \mathcal{A}$  such that  $A + B + C = 0$ , and an integer  $n \geq 1$ , there exist matrices  $A_n, B_n, C_n \in M_n(\mathbb{C})$  such that  $\lambda_{A_n} = \lambda_A^{(n)}, \lambda_{B_n} = \lambda_B^{(n)}, \lambda_{C_n} = \lambda_C^{(n)}$ , and  $A_n + B_n + C_n = 0$ .*

The proof of the relevant inequalities relies, as in finite dimensions, on finding projections with prescribed intersection properties. In order to state our main result in this direction we need a more precise description of the Horn inequalities. Assume that the subsets  $I = \{i_1 < i_2 < \dots < i_r\}$ ,  $J = \{j_1 < j_2 < \dots < j_r\}$ , and  $K = \{k_1 < k_2 < \dots < k_r\}$  of  $\{1, 2, \dots, n\}$  satisfy the identity

$$\sum_{\ell=1}^r [(i_\ell - \ell) + (j_\ell - \ell) + (k_\ell - \ell)] = 2r(n - r).$$

One associates to these sets a nonnegative integer  $c_{IJK}$ , called the Littlewood-Richardson coefficient. The sets  $I, J, K$  yield an eigenvalue inequality in Horn's conjecture if  $c_{IJK} \neq 0$ . Moreover, as shown by P. Belkale [1], the inequalities corresponding with  $c_{IJK} > 1$  are in fact redundant. Thus, the preceding theorem follows from the next result.

**Theorem 0.3.** *Given a factor  $\mathcal{A}$  of type  $II_1$ , selfadjoint elements  $A, B, C \in \mathcal{A}$  such that  $A + B + C = 0$ , an integer  $n \geq 1$ , and sets  $I, J, K \subset \{1, 2, \dots, n\}$  such that  $c_{IJK} = 1$ , we have*

$$\sum_{i \in I} \lambda_A^{(n)}(i) + \sum_{j \in J} \lambda_B^{(n)}(j) + \sum_{k \in K} \lambda_C^{(n)}(k) \leq 0.$$

This result follows from the existence of projections satisfying specific intersection requirements. Before stating our result in this direction, we need to specify a notion of genericity. Fix a finite factor  $\mathcal{A}$  with trace normalized so that  $\tau(1) = n$ . We will deal with flags of projections with integer dimensions, i.e., with collections

$$\mathcal{E} = \{0 = E_0 < E_1 < \dots < E_n = 1\}$$

of orthogonal projections in  $\mathcal{A}$  such that  $\tau(E_j) = j$  for all  $j$ . Given such a flag and a unitary operator  $U \in \mathcal{A}$ , the projections  $UE_jU^* = \{UE_jU^* : 0 \leq j \leq n\}$  form another flag. In fact, all flags with integer dimensions are obtained this way. A statement about a collection of three flags  $\{\mathcal{E}, \mathcal{F}, \mathcal{G}\}$  will be said to hold generically (or for generic flags) if it holds for the flags  $\{U\mathcal{E}U^*, V\mathcal{F}V^*, W\mathcal{G}W^*\}$  with  $(U, V, W)$  in a norm-dense open subset of  $\mathcal{U}(\mathcal{A})^3$  (where  $\mathcal{U}(\mathcal{A})$  denotes the group of unitaries in  $\mathcal{A}$ ). In finite dimensions, this set of unitaries can usually be taken to be Zariski open.

Note that flags with integer dimensions always exist if  $\mathcal{A}$  is of type  $II_1$ . If  $\mathcal{A} = M_m(\mathbb{C})$ , such flags only exist when  $n$  divides  $m$ .

One final piece of notation. Given variables  $\{e_j, f_j, g_j : 1 \leq j \leq n\}$ , we consider the free lattice  $L = L(\{e_j, f_j, g_j : 1 \leq j \leq n\})$ . This is simply the smallest collection which contains the given variables, and has the property that, given  $p, q \in L$ , the expressions  $(p) \wedge (q)$  and  $(p) \vee (q)$  also belong to  $L$ . We refer to the elements of  $L$  as *lattice polynomials*. If  $p$  is a lattice polynomial and  $\{E_j, F_j, G_j : 1 \leq j \leq n\}$  is a collection of orthogonal projections in a factor  $\mathcal{A}$ , we can substitute projections for the variables of  $p$  to obtain a new projection  $p(\{E_j, F_j, G_j : 1 \leq j \leq n\})$ . The lattice operations are interpreted as usual:  $P \vee Q$  is the projection onto the closed linear span of the ranges of  $P$  and  $Q$ , and  $P \wedge Q$  is the projection onto the intersection of the ranges of  $P$  and  $Q$ . Note that we did not impose any algebraic relations on  $L$ . When we work with flags, we can always reduce lattice polynomials using the relations  $e_j \wedge e_k = e_{\min\{j,k\}}$  and  $e_j \vee e_k = e_{\max\{j,k\}}$ . Further manipulations are possible because the lattice of projections in a finite factor is modular, i.e.  $(P \vee Q) \wedge R = P \vee (Q \wedge R)$  provided that  $P \leq R$ .

As in finite dimensions, the Horn inequalities follow from the intersection result below. Given a flag  $\mathcal{E} = (E_j)_{j=0}^n \subset \mathcal{A}$  such that  $\tau(E_j) = j$ , and a set  $I = \{i_1 < i_2 < \dots < i_r\} \subset \{1, 2, \dots, n\}$ , we denote by  $S(\mathcal{E}, I)$  the collection of projections  $P \in \mathcal{A}$  satisfying  $\tau(P) = r$  and

$$\tau(P \wedge E_{i_\ell}) \geq \ell, \quad \ell = 1, 2, \dots, r.$$

**Theorem 0.4.** *Given subsets  $I, J, K \subset \{1, 2, \dots, n\}$  with cardinality  $r$ , and with the property that  $c_{IJK} = 1$ , a finite factor  $\mathcal{A}$  with  $\tau(1) = n$ , and arbitrary flags  $\mathcal{E} = (E_j)_{j=0}^n$ ,  $\mathcal{F} = (F_j)_{j=0}^n$ ,  $\mathcal{G} = (G_j)_{j=0}^n$  such that  $\tau(E_j) = \tau(F_j) = \tau(G_j) = j$ , the intersection*

$$S(\mathcal{E}, I) \cap S(\mathcal{F}, J) \cap S(\mathcal{G}, K)$$

*is not empty.*

For generic flags, more is true.

**Theorem 0.5.** *Given subsets  $I, J, K \subset \{1, 2, \dots, n\}$  with cardinality  $r$ , and with the property that  $c_{IJK} = 1$ , there exists a lattice polynomial  $p \in L(\{e_j, f_j, g_j : 0 \leq j \leq n\})$  with the following property: for any finite factor  $\mathcal{A}$  with  $\tau(1) = n$ , and for generic flags  $\mathcal{E} = (E_j)_{j=0}^n$ ,  $\mathcal{F} = (F_j)_{j=0}^n$ ,  $\mathcal{G} = (G_j)_{j=0}^n$  such that  $\tau(E_j) = \tau(F_j) = \tau(G_j) = j$ , the projection  $P = p(\mathcal{E}, \mathcal{F}, \mathcal{G})$  has trace  $\tau(P) = r$  and, in addition*

$$\tau(P \wedge E_i) = \tau(P \wedge F_j) = \tau(P \wedge G_k) = \ell$$

*when  $i_\ell \leq i < i_{\ell+1}, j_\ell \leq j < j_{\ell+1}, k_\ell \leq k < k_{\ell+1}$  and  $\ell = 0, 1, \dots, r$ , where  $i_0 = j_0 = k_0 = 0$  and  $i_{r+1} = j_{r+1} = k_{r+1} = n + 1$ .*

When  $\mathcal{A} = M_n(\mathbb{C})$ , the existence and generic uniqueness of a projection  $P$  satisfying the trace conditions in the statement is well-known. In fact  $c_{IJK}$  serves as an algebraic way to count these projections. Our argument works equally well for linear subspaces of  $\mathbb{F}^n$  for any field  $\mathbb{F}$  (except that orthogonal complements  $1 - P$  must be replaced by annihilators in the dual). The following result is generally false when  $c_{IJK} > 1$ .

**Theorem 0.6.** *Fix a field  $\mathbb{F}$ , and complete flags  $\mathcal{E} = (E_j)_{j=0}^n$ ,  $\mathcal{F} = (F_j)_{j=0}^n$ ,  $\mathcal{G} = (G_j)_{j=0}^n$  of subspaces in  $\mathbb{F}^n$ . Given subsets  $I, J, K \subset \{1, 2, \dots, n\}$  with cardinality  $r$ , and with the property that  $c_{IJK} = 1$ , there exists a subspace  $M \subset \mathbb{F}^n$  such that  $\dim M = r$ , and*

$$\dim(M \cap E_{i_\ell}) \geq \ell, \quad \dim(M \cap F_{j_\ell}) \geq \ell, \quad \dim(M \cap G_{k_\ell}) \geq \ell$$

*for  $\ell = 1, 2, \dots, r$ .*

The search for projections  $P$  satisfying the conclusion of Theorem 0.4 is much more difficult when  $c_{IJK} > 1$ . One of the simplest cases of this problem is equivalent to the invariant subspace problem relative to a  $\text{II}_1$  factor  $\mathcal{A}$ ; this case was first discussed in [9] where an approximate solution is found. The relative invariant subspace problem remains open but there was spectacular progress in the work of U. Haagerup and H. Schultz [17, 18].

The function  $\lambda_{\mathcal{A}}$  can be defined more generally for a selfadjoint element of a von Neumann algebra  $\mathcal{A}$  endowed with a faithful, normal trace  $\tau$ . The inequalities in Theorem 0.3 are in fact true in this more general context. Rather than prove this fact directly, we will embed any such von Neumann algebra in a  $\text{II}_1$  factor, in such a way that the trace is preserved. An alternative proof can be obtained using von Neumann's reduction theory.

The remainder of this paper is organized as follows. In Section 1 we describe an enumeration of the sets  $I, J, K$  with  $c_{IJK} > 0$  in terms of a class of measures on the plane. This enumeration is essentially the one indicated in [22]; the measures we use can be viewed as the second derivatives of hives, or the first derivatives of honeycombs. (The fact that honeycombs enumerate Schubert intersection problems is proved in a direct way in the appendix of [8]; see also Tao's 'proof without words' illustrated in [26].) We also describe the duality observed in [22, Remark 2 on p. 42], realized by inflation to a puzzle, and \*-deflation to a dual measure. In Section 2, we use then the puzzle characterization of rigidity from [22] to formulate the condition  $c_{IJK} = 1$  in terms of the support of the corresponding measure  $m$ . This result may be viewed as the  $N = 0$  version of [22, Lemma 8]. This characterization is used in Section 3 to show that a measure  $m$  corresponding to sets with  $c_{IJK} = 1$  (also called a rigid measure) can be written uniquely as a sum  $m_1 + m_2 + \dots + m_p$  of extremal measures, and to introduce an order relation ' $\prec$ ' on the set  $\{m_j :$

$1 \leq j \leq p\}$ . In Section 4 we provide an extension of the concept of clockwise overlay from [22], and show that ‘ $\prec$ ’ provides examples of clockwise overlays. The main results are proved in Section 5. The most important observation is that general Schubert intersection problems can be reduced to problems corresponding to extremal measures. The order relation is essential here as the minimal measures  $m_j$  (relative to ‘ $\prec$ ’) must be considered first. A problem corresponding to an extremal measure has then a dual form (obtained by taking orthogonal complements) which is no longer extremal, except for essentially one trivial example. Section 6 contains a number of illustrations of this reduction procedure, including explicit expressions for the corresponding lattice polynomials which yield the solution for generic flags. In Section 7 we describe a particular intersection problem which is equivalent to the invariant subspace problem relative to a  $\mathrm{II}_1$  factor. In Section 8 we embed any algebra with a trace in a factor of type  $\mathrm{II}_1$ , and we show that projections can be moved to general position by letting them evolve according to free unitary Brownian motion.

There has been quite a bit of recent work on the geometry and intersection of Schubert cells. Belkale [2] shows that the inductive structure of the intersection ring of the Grassmannians can be justified geometrically. R. Vakil [26] provides an approach to the structure of this ring by a process of flag degenerations. He also indicates [27, 26] that this can be used in order to solve effectively all Schubert intersection problems, at least for generic flags. More precisely, [27, Remark 2.10] suggests that these solutions can be found, after an appropriate parametrization, by an application of the implicit function theorem which can be made numerically effective. These methods would apply to arbitrary values of  $c_{IJK}$ , and it is not clear that they would yield the formulas of Theorem 0.5 when  $c_{IJK} = 1$ . The method of flag degeneration of [26] depends essentially on finite dimensionality. A prospective analogue in a  $\mathrm{II}_1$  factor would require a checkerboard with a continuum of squares, and would be played with a continuum of pieces. This kind of game is difficult to organize, as illustrated for instance by the cumbersome argument used in [3]. The result proved with so much labor in that paper is deduced very simply from our current methods, as shown in Section 6 below.

## 1. HORN INEQUALITIES AND MEASURES

Fix integers  $1 \leq r \leq n$ , and subsets  $I, J, K \subset \{1, 2, \dots, n\}$  of cardinality  $r$ . We will find it useful on occasion to view the set  $I$  as an increasing function  $I : \{1, 2, \dots, r\} \rightarrow \{1, 2, \dots, n\}$ , i.e.,  $I = \{I(1) < I(2) < \dots < I(r)\}$ . The results of [22] show that we have  $c_{IJK} > 0$  if and only if there exist selfadjoint matrices  $X, Y, Z \in M_r(\mathbb{C})$  such that  $X + Y + Z = 2(n - r)1_{\mathbb{C}^r}$  and

$$\lambda_X(r + 1 - \ell) = I(\ell) - \ell, \lambda_Y(r + 1 - \ell) = J(\ell) - \ell, \lambda_Z(r + 1 - \ell) = K(\ell) - \ell$$

for  $\ell = 1, 2, \dots, r$ . Thus, as conjectured by Horn, such sets can be described inductively, using Horn inequalities with fewer terms. In other words, we have  $c_{IJK} > 0$  if and only if

$$\sum_{\ell=1}^r [(I(\ell) - \ell) + (J(\ell) - \ell) + (K(\ell) - \ell)] = 2r(n - r),$$


and

$$\sum_{\ell=1}^s [(I(I'(\ell)) - I'(\ell)) + (J(J'(\ell)) - J'(\ell)) + (K(K'(\ell)) - K'(\ell))] \geq 2s(n-r)$$

whenever  $s \in \{1, 2, \dots, r-1\}$  and  $I', J', K' \subset \{1, 2, \dots, r\}$  are sets of cardinality  $s$  such that  $c_{I'J'K'} > 0$ . The last inequality can also be written as

$$\sum_{\ell=1}^s [(I(I'(\ell)) - \ell) + (J(J'(\ell)) - \ell) + (K(K'(\ell)) - \ell)] \geq 2s(n-s).$$


The numbers  $c_{IJK}$  can be calculated using the Littlewood-Richardson rule which we discuss next. We use the form of the rule described in [22], so we need first to describe a set of measures on the plane. Begin by choosing three unit vectors  $u, v, w$  in the plane such that  $u + v + w = 0$ .



Consider the *lattice points*  $iu + jv$  with integer  $i, j$ . A segment joining two nearest lattice points will be called a *small edge*. We are interested in positive measures  $m$  which are supported by a union of small edges, whose restriction to each small edge is a multiple of linear measure, and which satisfy the balance condition (called zero tension in [22])

$$(1.1) \quad m(AB) - m(AB') = m(AC) - m(AC') = m(AD) - m(AD')$$

whenever  $A$  is a lattice point and the lattice points  $B, C', D, B', C, D'$  are in cyclic order around  $A$ .




If  $e$  is a small edge, the value  $m(e)$  is equal to the density of  $m$  relative to linear measure on that edge.

Fix now an integer  $r \geq 1$ , and denote by  $\triangle_r$  the (closed) triangle with vertices  $0, ru$ , and  $ru + rv = -rw$ . We will use the notation  $A_j = ju$ ,  $B_j = ru + jv$ , and  $C_j = (r-j)w$  for the lattice points on the boundary of  $\triangle_r$ . We also set

$$X_j = A_j + w, Y_j = B_j + u, Z_j = C_j + v$$

for  $j = 0, 1, 2, \dots, r+1$ . The following picture represents  $\triangle_5$  and the points just defined; the labels are placed on the left.



Given a measure  $m$ , a *branch point* is a lattice point incident to at least three edges in the support of  $m$ . We will only consider measures with at least one branch point. This excludes measures whose support consists of one or more parallel lines. We will denote by  $\mathcal{M}_r$  the collection of all measures  $m$  satisfying the balance condition above, whose branch points are contained in  $\triangle_r$ , and such that

$$m(A_j X_{j+1}) = m(B_j Y_{j+1}) = m(C_j Z_{j+1}) = 0, \quad j = 0, 1, \dots, r.$$

Analogously  $\mathcal{M}_r^*$  consists of measures  $m$  whose branch points are contained in  $\mathcal{M}_r$ , and such that

$$m(A_j X_j) = m(B_j Y_j) = m(C_j Z_j) = 0, \quad j = 0, 1, \dots, r.$$

Clearly,  $\mathcal{M}_r^*$  can be obtained from  $\mathcal{M}_r$  by reflection relative to one of the angle bisectors of  $\triangle_r$ .

Given a measure  $m \in \mathcal{M}_r$ , we define its *weight*  $\omega(m) \in \mathbb{R}_+$  to be

$$\omega(m) = \sum_{j=0}^r m(A_j X_j) = \sum_{j=0}^r m(B_j Y_j) = \sum_{j=0}^r m(C_j Z_j)$$

and its *boundary*  $\partial m = (\alpha, \beta, \gamma) \in (\mathbb{R}^r)^3$ , where

$$\alpha_\ell = \sum_{j=0}^{\ell-1} m(A_j X_j), \beta_\ell = \sum_{j=0}^{\ell-1} m(B_j Y_j), \gamma_\ell = \sum_{j=0}^{\ell-1} m(C_j Z_j), \quad \ell = 1, 2, \dots, r.$$

The equality of the three sums giving  $\omega(m)$  is an easy consequence of the balance condition.

The results of [21, 22] imply that the sets  $I, J, K \subset \{1, 2, \dots, n\}$  of cardinality  $r$  satisfy  $c_{IJK} > 0$  if and only if there exists a measure  $m \in \mathcal{M}_r$  with weight  $\omega(m) = n - r$ , and with boundary  $\partial m = (\alpha, \beta, \gamma)$  such that

$$\alpha_\ell = I(\ell) - \ell, \beta_\ell = J(\ell) - \ell, \gamma_\ell = K(\ell) - \ell, \quad \ell = 1, 2, \dots, r.$$

The number  $c_{IJK}$  is equal to the number of measures in  $\mathcal{M}_r$  satisfying these conditions, and with integer densities on all edges. Moreover, as shown in [22], if  $c_{IJK} = 1$ , there is only one measure  $m$  satisfying these conditions, and its densities must naturally be integers. In general, we will say that a measure  $m \in \mathcal{M}_r$  is *rigid* if it is entirely determined by its weight and boundary.



We will also use the version of these results in terms of  $\mathcal{M}_r^*$ , so we define for  $m \in \mathcal{M}_r^*$  the weight


$$\omega(m) = \sum_{j=0}^r m(A_j X_{j+1}) = \sum_{j=0}^r m(B_j Y_{j+1}) = \sum_{j=0}^r m(C_j Z_{j+1})$$

and boundary  $\partial m = (\alpha, \beta, \gamma)$ , where

$$\alpha_\ell = \sum_{j=r+1-\ell}^r m(A_j X_{j+1}), \beta_\ell = \sum_{j=r+1-\ell}^r m(B_j Y_{j+1}), \gamma_\ell = \sum_{j=r+1-\ell}^r m(C_j Z_{j+1})$$

for  $\ell = 1, 2, \dots, r$ .

Measures in  $\mathcal{M}_r$  or  $\mathcal{M}_r^*$  are entirely determined by their restrictions to  $\Delta_r$  and, when the corners of  $\Delta_r$  are not branch points, even by their restrictions to the interior of  $\Delta_r$ . Indeed, the lack of branch points outside  $\Delta_r$  implies that the densities are constant on the half-lines starting with  $A_j X_j, B_j Y_j$  and  $C_j Z_j$ . Note that a restriction  $m|_{\Delta_r}$  with  $m \in \mathcal{M}_r$  is not generally of the form  $m'|_{\Delta_r}$  for some  $m' \in \mathcal{M}_r^*$ . The first picture below represents  $\Delta_r$  (dotted lines), and the support (solid lines) of a measure in  $\Delta_r$ . The second one represents the support of a measure in  $\mathcal{M}_r^*$ .



To conclude this section, we establish a connection between measures in  $\mathcal{M}_r$  and the honeycombs of [21]. A *honeycomb* is a function  $h$  defined on the set of small edges contained in  $\Delta_r$  satisfying the following two properties:

- (i) If  $ABC$  is a small triangle contained in  $\Delta_r$ , we have  $h(AB) + h(AC) + h(BC) = 0$ .
- (ii) If  $A, B, C, D$  are lattice vertices in  $\Delta_r$  such that  $B = A + u$ ,  $C = A - v$ ,  $D = A + w$  (or  $B = A + v$ ,  $C = A - w$ ,  $D = A + u$ , or  $B = A + w$ ,  $C = A - u$ ,  $D = A + v$ ), then

$$h(AB) - h(CD) = h(BC) - h(AD) \geq 0.$$

The reason for the term honeycomb is not visible in our definition. One can associate to each small triangle  $ABC \subset \Delta_r$  the point  $(h(AB), h(BC), h(AC))$  in the plane  $\{(x, y, z) \in \mathbb{R}^3 : x + y + z = 0\}$ . These points form the vertices of a graph which looks like a honeycomb if it is not too degenerate (cf. [21]).

The following result will be required for our discussion of the Horn inequalities in Section 4.

**Lemma 1.1.** *Let  $m \in \mathcal{M}_r$  be a measure with weight  $\omega$  and  $\partial m = (\alpha, \beta, \gamma)$ . There exists a honeycomb  $h$  with the following properties.*

- (1)  $h(A_{\ell-1}A_\ell) = \alpha_\ell - 2\omega/3$ ,  $h(B_{\ell-1}B_\ell) = \beta_\ell - 2\omega/3$ ,  $h(C_{\ell-1}C_\ell) = \gamma_\ell - 2\omega/3$  for  $\ell = 1, 2, \dots, r$ .
- (2) If  $B = A + u$ ,  $C = A - v$ ,  $D = A + w$  (or  $B = A + v$ ,  $C = A - w$ ,  $D = A + u$ , or  $B = A + w$ ,  $C = A - u$ ,  $D = A + v$ ) then  $h(AB) - h(CD) = m(AC)$ .

*Proof.* This is routine. Condition (2) allows us to calculate all the values of  $h$  starting from the boundary of  $\Delta_r$ . To verify (ii) one must use the balance condition for measures in  $\mathcal{M}_r$ .  $\square$

## 2. INFLATION, DUALITY, AND RIGIDITY


Let  $\mathcal{A}$  be a finite factor with trace normalized so that  $\tau(1) = n$ , and let  $\mathcal{E} = \{E_\ell : \ell = 0, 1, \dots, n\}$  be a flag so that  $\tau(E_\ell) = \ell$  for all  $\ell$ . Fix also a set  $I \subset \{1, 2, \dots, n\}$  of cardinality  $r$  and a projection  $P \in \mathcal{A}$  with  $\tau(P) = r$ . We have  $P \in S(\mathcal{E}, I)$ , i.e.  $\tau(P \wedge E_{I(\ell)}) \geq \ell$  for  $\ell = 1, 2, \dots, r$ , if and only if  $\tau(P \wedge E_\ell) \geq \varphi_I(\ell)$  for  $\ell = 0, 1, \dots, n$ , where

$$\varphi_I(\ell) = p \text{ for } I(p) \leq \ell < I(p+1)$$

for  $p = 0, 1, \dots, r$ , and  $I(0) = 1, I(r+1) = n+1$ . With the notation  $P^\perp = 1 - P$ , these conditions imply

$$\begin{aligned} \tau(P^\perp \wedge E_\ell^\perp) &= n - \tau(P \vee E_\ell) \\ &= n - \tau(P) - \tau(E_\ell) + \tau(P \wedge E_\ell) \\ &\geq n - r - \ell + \varphi_I(\ell). \end{aligned}$$


This implies that  $P \in S(\mathcal{E}, I)$  if and only if  $P^\perp \in S(\mathcal{E}^\perp, I^*)$ , where  $\mathcal{E}^\perp = \{E_{n-\ell}^\perp : \ell = 0, 1, \dots, n\}$ , and  $I^* = \{n+1-i : i \notin I\}$ . In general, we will have  $c_{I^*J^*K^*} = c_{IJK}$ , and this equality is realized by a duality considered in [22]. More precisely, assume that  $m \in \mathcal{M}_r$ . We define the *inflation* of  $m$  as follows. Cut  $\Delta_r$  along the edges in the support of  $m$  to obtain a collection of (white) puzzle pieces, and translate these pieces away from each other in the following way: the parallelogram formed by the two translates of a side  $AB$  of a white puzzle piece has two sides of length equal to the density of  $m$  on  $AB$  and  $60^\circ$  clockwise from  $AB$ . The original puzzle pieces and these parallelograms fit together, and leave a space corresponding to each branch point in the support of  $m$ . Here is an illustration of the process; the thinner lines in the support of the measure have density one, and the thicker ones density 2. The original pieces of the triangle  $\Delta_r$  are white, the added parallelogram pieces are dark gray, and the branch points become light gray pieces. Each light gray piece has as many sides as there are branches at the original branch point (counting the branches outside  $\Delta_r$ , which are not represented in this figure, though their number and densities are dictated by the balance condition, and the fact that  $m$  belongs to  $\mathcal{M}_r$ ).



The original triangle  $\Delta_r$  has been inflated to a triangle of size  $r + \omega(m)$ , and the decomposition of this triangle into white, gray, and light gray pieces is known as the *puzzle* associated to  $m$ . Each gray parallelogram has two *light gray sides*, i.e. sides bordering a light gray piece, and two *white sides*. The length of the light gray side equals the density of the white sides in the original support of  $m$  in  $\Delta_r$ . This process can be applied to the entire support of  $m$ , but we are only interested in its

effect on  $\Delta_r$ . (The white regions in the puzzle are called ‘zero regions’, and the light gray ones ‘one regions’ in [22]. We use in our drawings a color scheme different from the one used in [22].)

We can now apply a dual deflation, or *\*deflation*, to the puzzle of  $m$  as follows: discard all the white pieces, and shrink the gray parallelograms by reducing their white sides to points. The segments obtained this way are assigned densities equal to the lengths of the white sides of the corresponding parallelograms. In the picture below, the shrunk parallelograms are represented as solid lines.




The result of this deflation is a triangle with sides  $\omega(m)$ , endowed with a measure supported by the solid lines which will be denoted  $m^*$ . The support of  $m^*$  can be obtained directly from the support of  $m$  as follows: take every edge of a white puzzle piece, rotate it  $60^\circ$  clockwise, and change its length to the density of  $m$  on the original edge. The new segments must now be translated so that the edges originating from the sides of a white puzzle piece become concurrent. Thus, the dual picture depends primarily on the combinatorial structure of the support of  $m$ . More precisely, let us say that the measures  $m \in \mathcal{M}_r$  and  $m' \in \mathcal{M}_{r'}$  are *homologous* if there is a bijection between the edges determined by the support of  $m$  and the edges determined by the support of  $m'$  such that corresponding edges are parallel, and concurrent edges correspond with concurrent edges (the concurrence point being precisely the one dictated by the correspondence of the edges). Then  $m$  and  $m'$  are homologous if and only if  $m^*$  and  $m'^*$  are homologous. For instance, measures in  $\mathcal{M}_r$  that have the same support are homologous.

Assume now that the measure  $m \in \mathcal{M}_r$  has integer densities, and  $I, J, K$  are the corresponding sets in  $\{1, 2, \dots, n = r + \omega(m)\}$ . Then the triangle obtained by inflating  $m$  can be identified with  $\Delta_n$ . Under this identification the small edges  $A_{i-1}A_i$  are either white (if they border a white piece, or they belong to a white edge of a gray parallelogram) or light gray. It is easy to see that the white small edges are precisely  $A_{i_\ell-1}A_{i_\ell}$  for  $\ell = 1, 2, \dots, r$ , and therefore the light gray edges correspond with the complement of  $I$ . Furthermore, the light gray triangle obtained by *\*deflation* can be identified with  $\Delta_{n-r}$ , and  $m^* \in \mathcal{M}_{n-r}^*$  is a measure satisfying  $\omega(m^*) = r$ . This measure determines subsets of  $\{1, 2, \dots, n\}$  which are precisely  $I^*, J^*, K^*$ . This observation gives a bijective proof of the equality  $c_{IJK} = c_{I^*J^*K^*}$ .

The passage from  $m$  to  $m^*$  can be reversed by applying *\*inflation* to  $m^*$ , and then applying deflation to the resulting puzzle.


Another important application of the inflation process is a characterization of rigidity. Orient the edges of the gray parallelograms in a puzzle so that they point away from the acute angles. Some of the border edges do not have a neighboring gray parallelogram and will not be oriented.



It was shown in [22] that a measure  $m$  is rigid if and only if the associated directed graph contains no gentle loops, i.e., loops which never turn more than  $60^\circ$ . Note that the number and relative position of the puzzle pieces depends only on the support of the measure  $m$ . The following result follows immediately.

**Proposition 2.1.** *Let  $m_1, m_2 \in \mathcal{M}_r$  be such that the support of  $m_1$  is contained in the support of  $m_2$ . If  $m_2$  is rigid then  $m_1$  is rigid as well.*

It is also easy to see that the support of a rigid measure does not contain six edges which meet at the same point. Indeed, the inflation reveals immediately a gentle loop.



We will need to characterize rigidity in terms of the support of the original measure. Let  $A_1 A_2 \cdots A_k A_1$  be a loop consisting of small edges  $A_j A_{j+1}$  contained in the support of a measure  $m \in \mathcal{M}_r$ . We will say that this loop is *evil* if each three consecutive points  $A_{j-1} A_j A_{j+1} = ABC$  form an *evil turn*, i.e. one of the following situations occurs:

- (1)  $C = A$ , and the small edges  $BX, BY, BZ$  which are  $120^\circ, 180^\circ$ , and  $240^\circ$  clockwise from  $AB$  are in the support of  $m$ .
- (2)  $BC$  is  $120^\circ$  clockwise from  $AB$ .
- (3)  $C \neq A$  and  $A, B, C$  are collinear.
- (4)  $BC$  is  $120^\circ$  counterclockwise from  $AB$  and the edge  $BX$  which is  $120^\circ$  clockwise from  $AB$  is in the support of  $m$ .
- (5)  $BC$  is  $60^\circ$  counterclockwise from  $AB$  and the edges  $BX, BY$  which are  $120^\circ$  and  $180^\circ$  clockwise from  $AB$  are in the support of  $m$ .


**Proposition 2.2.** *A measure  $m \in \mathcal{M}_r$  is rigid if and only if its support contains none of the following configurations:*

- (1) Six edges meeting in one lattice point;
- (2) An evil loop.


*Proof.* Assume first that  $m$  is not rigid, and consider a gentle loop of minimal length in its puzzle. Use  $a_i$  to denote white parallelogram sides in this loop, and  $b_i$  light gray parallelogram sides. The sides  $a_i, b_i$  may consist of several small edges. The gentle loop is of one of the following three forms:

- (1)  $a_1 a_2 a_3 a_4 a_5 a_6$ ,
- (2)  $b_1 b_2 b_3 b_4 b_5 b_6$ ,
- (3)  $a_1 a_2 \cdots a_{i_1} b_1 b_2 \cdots b_{j_1} a_{i_1+1} \cdots a_{i_2} b_{j_1+1} \cdots b_{j_2} \cdots a_{i_{p-1}+1} \cdots a_{i_p} b_{j_{p-1}+1} \cdots b_{j_p}$ ,  
with at most five consecutive  $a$  or  $b$  symbols.


In case (1), the loop runs counterclockwise around a white piece, and it deflates to a translation of itself which is obviously evil. In case (2), the loop deflates to a single point where six edges in the support of  $m$  meet. In case (3), the loop deflates to  $a'_1 a'_2 \cdots a'_{i_p}$ , where each  $a'_j$  is a translate of  $a_j$ . The turns in this loop are obtained by deflating a path of the form  $a_1 b_1 \cdots b_j a_2$  with  $0 \leq j \leq 5$ . The edges  $b_1 \cdots b_j$  run clockwise around a light gray piece which must have at most five edges because the gentle loop was taken to have minimal length. When  $j = 0$ , the edges  $a_1$  and  $a_2$  border the same white puzzle piece, and it is obvious that  $a'_1 a'_2$  is an evil turn. The remaining cases will be enumerated according to the number of edges in the light gray piece next to the edges  $b_j$ . When this piece is a triangle, we can only have  $j = 1$ , and the situation is illustrated below. The dashed line in the deflation indicates a portion of the support of  $m$ .




When the light gray piece is a parallelogram, we have  $j = 1$  or  $j = 2$ . The three possible deflations are as follows.




Next, the light gray piece may be a trapezoid, and  $1 \leq j \leq 3$ . For  $j = 1$  we have these four possibilities:




For  $j = 2, 3$  there are three more possibilities.




Finally, if the light gray piece is a pentagon, there are five situations when  $j = 1$ ,




four situations when  $j = 2$ ,



three situations when  $j = 3$ ,




and three more when  $j = 4, 5$ .



Of course, the last case does not occur in a minimal gentle loop.

Thus, in all situations, the deflated turns are evil, and therefore the support of  $m$  contains an evil loop. Conversely, if the support of  $m$  contains an evil loop, the above figures show that one can obtain a gentle loop in the puzzle of  $m$ .  $\square$


The following figure represents the support of a rigid measure in  $\mathcal{M}_8$ , along with a loop which may seem evil but is not evil when traversed in either direction.




### 3. EXTREMAL MEASURES AND SKELETONS

For fixed  $r \geq 1$ , the collection  $\mathcal{M}_r$  is a convex polyhedral cone. Recall that a measure  $m \in \mathcal{M}_r$  is *extremal* (or belongs to an extreme ray) if any measure  $m' \leq m$  is a positive multiple of  $m$ . The support of an extremal measure will be called a *skeleton*. Clearly, an extremal measure is entirely determined by its value on any small edge contained in its skeleton. Checking extremality is easily done by using the balance condition (1.1) at all the branch points of the support to see how the density propagates from one edge to the rest of the support.


In the following figure of a skeleton, the thicker edges must be assigned twice the density of the other edges. (This skeleton contains an evil loop, hence the measures it supports are not rigid.)




It is not always obvious when a collection of edges supports a nonzero measure in  $\mathcal{M}_r$ . The reader may find it amusing to verify that the following figure represents sets which do not support any nonzero measure.




We will be mostly interested in the supports of rigid extremal measures, which we will call rigid skeletons. When  $r = 1$ , there are only the three possible skeletons, all of them rigid, pictured below.




The following figure shows some rigid skeletons for  $r = 2, 3, 4, 5$ .



A greater variety of rigid skeletons is available for  $r = 6$ . In addition to larger versions (plus rotations and reflections) of the above skeletons, we have the ones in the next figure.



For larger  $r$ , rigid skeletons can be quite involved. We provide just one more example for  $r = 8$ . This skeleton has edges with densities 2 and 3 which we did not indicate.



An important consequence of the characterization of rigid measures in  $\mathcal{M}_r$  is the fact that such measures can be written uniquely as sums of extremal measures. Fix a rigid measure  $m \in \mathcal{M}_r$ , and let  $e = AB$  and  $f = BC$  be two distinct small edges. We will write  $e \rightarrow_m f$ , or simply  $e \rightarrow f$  when  $m$  is understood, if one of these two situations arises:

- (1)  $\angle ABC = 120^\circ$  and the edge opposite  $e$  at  $B$  has  $m$  measure equal to zero;
- (2)  $e$  and  $f$  are opposite, and one of the edges making an angle of  $60^\circ$  with  $f$  has  $m$  measure equal to zero.

Note that in both cases we may also have  $f \rightarrow e$ . The significance of this relation is that  $e \rightarrow f$  implies that  $m(e) \leq m(f)$ , with strict inequality unless  $f \rightarrow e$  as well. More precisely, if  $e \rightarrow f$  and we do not have  $f \rightarrow e$ , there is at least one edge



$g$  such that  $g \rightarrow f$  and the angle between  $e$  and  $g$  is  $60^\circ$ . In this case we have

$$(3.1) \quad m(f) = m(e) + m(g).$$

Indeed, if  $e$  and  $f$  are opposite, the edge opposite  $g$  must have  $m$  measure equal to zero.

A useful observation is that if  $XY \rightarrow YZ$  but  $YZ \not\rightarrow XY$ , then the edges  $YA, YB, YC$  which are  $120^\circ, 180^\circ, 240^\circ$  clockwise from  $YZ$  must be in the support of  $m$ . In other words,  $ZYZ$  is an evil turn.

The following result is a simple consequence of the fact that the measure  $m$  exists.

**Lemma 3.1.** *Assume that a sequence of edges  $e_1, e_2, \dots, e_n$  in the support of  $m$  is such that*

$$e_1 \rightarrow e_2 \rightarrow e_3 \rightarrow \dots \rightarrow e_n \rightarrow e_1.$$

*Then we also have*

$$e_n \rightarrow e_{n-1} \rightarrow \dots \rightarrow e_1 \rightarrow e_n.$$

*Proof.* Indeed, if one of the arrows cannot be reversed, then  $m(e_1) < m(e_1)$ .  $\square$

We can therefore define a preorder relation on the set of small edges as follows. Given two small edges  $e, f$ , we write  $e \Rightarrow f$  if either  $e = f$ , or there exist edges  $e_1, e_2, \dots, e_n$  such that

$$e = e_1 \rightarrow e_2 \rightarrow \dots \rightarrow e_n = f.$$

In this case, we will say that  $f$  is a *descendant* of  $e$  and  $e$  is an *ancestor* of  $f$ . Two edges are *equivalent*,  $e \Leftrightarrow f$ , if  $e \Rightarrow f$  and  $f \Rightarrow e$ . The relation of descentance becomes an order relation on the equivalence classes of small edges. An edge  $e$  will be called a *root* if  $m(e) \neq 0$  and  $e$  belongs to a minimal class relative to descentance. Clearly, every edge in the support of  $m$  is a descendant of at least one root.

If  $m(e) \neq 0$  and  $e \Rightarrow f$ , then there exists a path  $A_0 A_1 \dots A_k$  in the support of  $m$  such that  $A_{j-1} A_j \rightarrow A_j A_{j+1}$  for all  $j$ ,  $A_0 A_1 = e$ , and  $A_{k-1} A_k = f$ . Paths of this form will be referred to as *descentance paths* from  $e$  to  $f$ . All the turns  $A_{j-1} A_j A_{j+1}$  and  $A_{j+1} A_j A_{j-1}$  in a descentance path are evil.

**Lemma 3.2.** *Assume that  $e$  and  $f$  are in the support of a rigid measure  $m \in \mathcal{M}_r$ , and  $A_0 A_1 \dots A_k$  and  $B_0 B_1 \dots B_\ell$  are two descentance paths from  $e$  to  $f$ . Then  $A_{k-1} = B_{\ell-1}$  and  $A_k = B_\ell$ .*

*Proof.* Assume to the contrary that  $A_{k-1} \neq B_{\ell-1}$ . There are two cases to consider, according to whether  $A_0 = B_1$  or  $A_0 = B_0$ . In the first case, the loop

$$A_0 A_1 \dots A_{k-1} B_{\ell-1} B_{\ell-2} \dots B_1$$

is evil, contradicting the rigidity of  $m$ . In the second case, there is a first index  $p$  such that  $A_{p+1} \neq B_{p+1}$ . Then the loop

$$A_p A_{p+1} \dots A_{k-1} B_{\ell-1} B_{\ell-2} \dots B_p$$

is evil, yielding again a contradiction.  $\square$

**Lemma 3.3.** *Let  $e$  and  $e'$  be inequivalent root edges in the support of a rigid measure  $m \in \mathcal{M}_r$ , and let  $f$  be an edge which is a descendant of both  $e$  and  $e'$ . Consider a descentance path  $A_0 A_1 \dots A_k$  from  $e$  to  $f$ , and a descentance path  $B_0 B_1 \dots B_\ell$  from  $e'$  to  $f$ . Then  $A_{k-1} = B_{\ell-1}$  and  $A_k = B_\ell$ .*

*Proof.* Assume to the contrary that  $A_k = B_{\ell-1}$ . The edge  $f$  is not equivalent to either  $e$  or  $e'$ . Therefore there exist indices  $p, q$  such that  $A_p A_{p+1} \not\rightarrow A_{p-1} A_p$  and  $B_q B_{q+1} \not\rightarrow B_{q-1} B_q$ . It follows that

$$A_p A_{p+1} \cdots A_{k-1} B_{\ell-1} B_{\ell-2} \cdots B_q B_{q-1} \cdots B_{\ell-2} B_{\ell-1} A_{k-1} \cdots A_{p+1} A_p$$

is an evil loop, contradicting rigidity.  $\square$

These lemmas show that all the non-root edges in the support of a rigid measure  $m$  can be given an orientation. More precisely, given a relation  $e \Rightarrow f$  with  $e$  a root edge, choose a descendance path  $A_0 A_1 \cdots A_k$  from  $e$  to  $f$ , and assign  $f$  the orientation  $A_{k-1} A_k$ . This will be called the *orientation of  $f$  away from the root edges*. Any common edge of two skeletons in the support of  $m$  can be oriented away from the root edges; indeed, such an edge is not a root edge. In the proofs of the next two results, we will be concerned with the descendants of a fixed root edge  $e$ , and it will be convenient to orient the other root edges equivalent to  $e$  away from  $e$ . The edge  $e$  can be oriented either way, as needed.

**Lemma 3.4.** *Fix a root edge in the support of a rigid measure  $m$ , and suppose that two descendants  $f = CX$  and  $g = DX$  have orientations pointing toward  $X$ . Then the turns  $CXD$  and  $DXC$  are not evil. In particular, the angle between  $f$  and  $g$  is  $60^\circ$ , and at least one of the edges  $C'X, D'X$  opposite  $f$  and  $g$  has  $m$  measure equal to zero.*

*Proof.* Assume to the contrary that either  $CXD$  or  $DXC$  is an evil turn. The assumed orientations imply that  $f \not\rightarrow g$  and  $g \not\rightarrow f$ . Since one of the edges incident to  $X$  must have measure zero, it follows  $f$  and  $g$  are not collinear. Moreover, the edges  $f' = XC'$  and  $g' = XD'$  opposite to  $f$  and  $g$ , respectively, must be in the support of  $m$ ; in the contrary case we would have  $f \rightarrow g$  or  $g \rightarrow f$  if the angle between  $f$  and  $g$  is  $120^\circ$ , or the turn  $CXD$  would not be evil if the angle is  $60^\circ$ . Let  $A_0 A_1 \cdots A_k$  and  $B_0 B_1 \cdots B_\ell$  be descendance paths from  $e$  to  $f$  and  $g$ , respectively. By assumption, we have  $A_{k-1} = C$ ,  $B_{\ell-1} = D$ , and  $A_k = B_\ell = X$ . If  $A_0 \neq B_0$ , then  $A_0 = B_1$ ,  $A_1 = B_0$ , and clearly

$$A_0 A_1 \cdots A_k B_{\ell-1} B_{\ell-2} \cdots B_1$$

or its reverse is an evil loop, contradicting the rigidity of  $m$ . Thus we must have  $A_0 = B_0$ . Let  $p$  be the largest integer such that  $A_j = B_j$  for  $j \leq p$ . If  $p < \min\{k, \ell\}$ , the loop

$$A_p A_{p+1} \cdots A_k B_{\ell-1} B_{\ell-2} \cdots B_p$$

or its reverse is evil. Indeed, since  $A_{p-1} A_p \rightarrow A_p A_{p+1}$  and  $A_{p-1} A_p \rightarrow B_p B_{p+1}$ , the turns  $A_{p+1} B_p B_{p+1}$  and  $B_{p+1} B_p A_{p+1}$  are both evil. We conclude that  $p = \min\{k, \ell\}$ . If  $p = k$ , it follows that the  $B_{k-1} B_k \cdots B_\ell$  is a descendance path from  $f$  to  $g$ . Since  $f \not\rightarrow g$ , we must have  $B_{k+1} = C'$ , and then the loop

$$B_k B_{k+1} \cdots B_\ell$$

or its reverse is obviously evil, leading to a contradiction. The case  $p = \ell$  similarly leads to a contradiction.  $\square$

**Theorem 3.5.** *Let  $m \in \mathcal{M}_r$  be a rigid measure, and  $e$  a root edge in the support of  $m$ . Then the collection of all descendants of  $e$  is a skeleton.*

*Proof.* Since  $m$  can be written as a sum of extremal measures, there exists an extremal measure  $m' \leq m$  such that  $m'(e) \neq 0$ . Since  $f \rightarrow_m g$  implies that  $f \rightarrow_{m'} g$ , the support of  $m'$  is a skeleton containing all the descendants of  $e$ . Therefore it will suffice to show that the descendants of  $e$  form the support of some measure in  $\mathcal{M}_r$ . We set  $\mu(e) = 1$ ,  $\mu(f) = 0$  if  $f$  is not a descendant of  $e$ , and for each descendant  $f \neq e$  of  $e$  we define  $\mu(f)$  to be the number of descendance paths from  $e$  to  $f$ . Clearly, no edge occurs twice in such a path; such an occurrence would imply the existence of an evil loop. Thus the number  $\mu(f)$  is finite. To conclude the proof, it suffices to show that  $\mu \in \mathcal{M}_r$ . The support of  $\mu$  is contained in the support of  $m$ . Therefore all the branch points are in  $\triangle_r$ , and

$$\mu(A_j X_{j+1}) = \mu(B_j Y_{j+1}) = \mu(C_j Z_{j+1}) = 0$$

for all  $j$ . It remains to verify the balance conditions. Consider a lattice point  $X$  in  $\triangle_r$ . If no descendant of  $e$  is incident to  $X$ , the six edges meeting at  $X$  have  $\mu$  measure zero, and the balance condition is trivial. Otherwise, the number of descendants of  $e$  incident to  $X$  can be 1, 2, 3, 4 or 5; the value 6 is excluded by the rigidity of  $m$ . We first exclude the case where this number is 1. Assume indeed that there is only one descendant incident to  $X$ , and let  $A_0 A_1 \cdots A_k$  be a descendance path from  $e$  with  $A_k = X$ . Since  $A_{k-1} A_k$  has no descendants of the form  $XY$ , it follows that the turn  $A_{k-1} A_k A_{k-1}$  is evil. On the other hand, since rigidity of  $m$  insures that one of the edges around  $X$  has measure zero, the edge  $A_{k-1} A_k$  is a strict descendant of some other edge  $XZ$ . In particular,  $A_{k-1} A_k$  is not a root edge, and therefore it is not equivalent to  $e$ . It follows that, for some  $p$ , we do not have  $A_p A_{p+1} \rightarrow A_{p-1} A_p$ , and this implies that the turn  $A_{p+1} A_p A_{p+1}$  is evil. Thus

$$A_p A_{p+1} \cdots A_{k-1} A_k A_{k-1} \cdots A_{p+1} A_p$$

is an evil loop, contrary to the rigidity of  $m$ .

Consider now the case when there are exactly two descendants of  $e$  incident to  $X$ , call them  $f$  and  $g$ . They cannot both point away from  $X$  since this would require the existence of a third descendant pointing toward  $X$ . They cannot both point toward  $X$ . Indeed, if this were the case, Lemma 3.4 insures that one of the edges opposite  $f$  and  $g$  has measure zero, and therefore  $f$  or  $g$  has another descendant pointing away from  $X$ . Thus we can assume that  $f$  points toward  $X$ , and  $g$  away from  $X$ , in which case we have  $f \rightarrow g$ . Then  $f$  and  $g$  must be collinear, and every descendance path for  $g$  passes through  $f$ . Thus  $\mu(f) = \mu(g)$ , which is the required balance condition.


Assume next that there are exactly three descendants incident to  $X$ . Two of them must be noncollinear and of the form  $WX \rightarrow XY$ , and the third descendant must be  $WX \rightarrow XZ$ , with the three edges forming  $120^\circ$  angles. Every descendance path for either  $XY$  or  $XZ$  passes through  $WX$ , showing that  $\mu(WX) = \mu(XY) = \mu(XZ)$ , and therefore satisfying the balance requirement at  $X$ .

Now, consider the case of exactly four descendants incident to  $X$ . If these four edges form two collinear pairs, then two of them must point towards  $X$ , and they will form an evil turn, contrary to Lemma 3.4. Therefore we can find among the four descendants two noncollinear edges  $WX \rightarrow XY$ , in which case we also have  $WX \rightarrow XZ$  with these three edges forming  $120^\circ$  angles. The fourth descendant is not collinear with  $WX$ , so it makes a  $60^\circ$  angle with  $WX$ . If it points away from  $X$ , it must be a descendant of the only incoming edge  $WX$ , and this is not possible. Therefore this fourth edge must be  $VX$  with  $VX \rightarrow XY$  or  $VX \rightarrow XZ$ . Assume

$VX \rightarrow XY$  for definiteness. In this case, all descendance paths for  $XZ$  pass through  $WX$ , so that  $\mu(XZ) = \mu(WX)$ . On the other hand, descendance paths for  $XY$  pass either through  $WX$  or through  $VX$ , showing that  $\mu(XY) = \mu(WX) + \mu(VX)$ . The balance requirement is again verified.

Finally, if 5 descendants of  $e$  are incident to  $X$ , then the sixth edge must have mass equal to zero, and it is impossible to orient the five edges so that every pair of incoming edges form a  $60^\circ$  angle, and every outgoing edge is a descendant of an incoming edge. Thus, this situation does not occur.  $\square$

The preceding proof shows that a rigid skeleton does not cross itself transversely. In other words, a rigid skeleton does not contain four edges meeting at the same point, such that they form two collinear pairs of edges. The following figure shows the possible ways (up to rotation) that the edges of a skeleton can meet, along with the possible orientations. In each case, a (or the) dotted edge must have density equal to zero.



The measure constructed in the above argument is the only measure supported by the descendants of  $e$  such that  $\mu(e) = 1$ . We will denote this measure  $\mu_e$ .

**Corollary 3.6.** *Let  $m \in \mathcal{M}_r$  be a rigid measure, and let  $e_1, e_2, \dots, e_k$  be a maximal collection of inequivalent root edges. Then we have*

$$m = \sum_{j=1}^k m(e_j) \mu_{e_j}.$$

*Proof.* Let us say that an edge  $f$  in the support of  $m$  has height  $\geq p \geq 2$  if there exists a descendance path  $A_0 A_1 \dots A_p$  from some root edge  $e$  to  $f$ . If  $f$  has height  $\geq p$ , we say that  $f$  has height equal to  $p$  if it does not have height  $\geq p+1$ . The requirement that  $m \in \mathcal{M}_r$  shows that the measure of any edge can be calculated in terms of the measures of edges of smaller height, as can be seen from (3.1). Therefore,  $m$  is entirely determined by the values  $m(e_j)$ ,  $j = 1, 2, \dots, k$ . On the other hand, the measure  $m' = \sum_{j=1}^k m(e_j) \mu_{e_j}$  has support contained in the support of  $m$ . Since  $m'(e_j) = m(e_j)$ , we conclude that  $m = m'$ .  $\square$

We mention one more useful property of rigid skeletons.

**Lemma 3.7.** *Let  $e$  and  $f$  be two edges in a rigid skeleton. There exists a path  $C_0 C_1 \dots C_p$  in this skeleton such that  $C_0 C_1 = e$ ,  $C_{p-1} C_p = f$ , and all the turns  $C_{j-1} C_j C_{j+1}$  and  $C_{j+1} C_j C_{j-1}$  are evil.*

*Proof.* The result is obvious if  $e$  is a root edge. If it is not, choose descendance paths  $A_0 A_1 \dots A_k$  from a root edge to  $e$ , and  $B_0 B_1 \dots B_\ell$  from the same root edge to  $f$ . If  $A_0 = B_1$ , the path

$$A_k A_{k-1} \dots A_1 B_1 B_2 \dots B_\ell$$


satisfies the requirements. If  $A_0 = B_0$  one chooses instead the path

$$A_k A_{k-1} \dots A_{r+1} B_r B_{r-1} \dots B_\ell,$$

where  $r$  is the first integer such that  $A_{r+1} \neq B_{r+1}$ . If no such integer exists, one of the paths is contained in the other. For instance,  $k < \ell$  and  $A_j = B_j$  for  $j \leq k$ . In this case the desired path is  $B_{k-1}B_k \cdots B_\ell$ .  $\square$


The path provided by this lemma is not generally a descent path.

We conclude this section by introducing an order relation on the set of skeletons contained in the support of a rigid measure  $m$ . Given two rigid skeletons  $S_1$  and  $S_2$ , we will write  $S_1 \prec_0 S_2$  if  $S_1$  has collinear edges  $AX, XB$  and  $S_2$  has collinear edges  $CX, XD$  such that  $XA$  is  $60^\circ$  clockwise from  $XC$ . The following figure shows the four possible configurations of  $S_1$  and  $S_2$  around the point  $X$ , up to rotation, and assuming that the two skeletons are contained in the support of a rigid measure. The edges in  $S_1 \setminus S_2$  are dashed, the edges in  $S_2 \setminus S_1$  are solid without arrows, and the common edges are oriented away from the root edges.



The four turns  $AXC, AXD, BXC$ , and  $BXD$  are evil.

Note that the point  $X$  could be on the boundary of  $\Delta_r$ , but not one of the three corner vertices. It is possible that  $S_1 \prec_0 S_2$  and  $S_2 \prec_0 S_1$ , as illustrated in the picture below (with  $S_1$  in dashed lines).



We will show that this does not occur when the skeletons are associated with a fixed rigid measure.

**Theorem 3.8.** *Fix a rigid measure  $m \in \mathcal{M}_r$  and an integer  $n \geq 1$ . There do not exist skeletons  $S_1, S_2, \dots, S_n$  contained in the support of  $m$  such that*

$$S_1 \prec_0 S_2 \prec_0 \cdots \prec_0 S_n \prec_0 S_1.$$


*Proof.* Assume to the contrary that such skeletons exist. Choose for each  $j$  collinear edges  $A_jX_j, X_jB_j$  in the support of  $S_j$  and collinear edges  $C_jX_j, X_jD_j$  in the support of  $S_{j+1}$  (with  $S_{n+1} = S_1$ ) such that  $XA_j$  is  $60^\circ$  clockwise from  $XC_j$ . The rigidity of  $m$  implies that one of the edges  $XY$  has measure zero. Therefore we must have either  $X_jA_j \rightarrow X_jB_j$  or  $X_jB_j \rightarrow X_jA_j$ . Label these two edges  $f_j$  and  $f'_j$  so that  $f_j \rightarrow f'_j$ , and note that  $f_j$  is not the descendant of any edge  $XY$ , except possibly  $f'_j$ . Analogously, denote the edges  $X_{j-1}C_{j-1}$  and  $X_{j-1}D_{j-1}$  by  $e_j$  and  $e'_j$  so that  $e'_j \rightarrow e_j$ . Since both  $e_j$  and  $f_j$  are contained in  $S_j$ , Lemma 3.7 provides a path with evil turns joining  $e_j$  and  $f_j$ . A moment's thought shows that this path either begins at  $X_{j-1}$ , or it begins with one of  $C_{j-1}X_{j-1}D_{j-1}, D_{j-1}X_{j-1}C_{j-1}$ . If

the second alternative holds, remove the first edge from the path. Performing the analogous operation at the other endpoint, we obtain a path  $\gamma_j$  with only evil turns which starts at  $X_{j-1}$  ends at  $X_j$  (with  $X_{j-1} = X_n$  if  $j = 0$ ), its first edge is one of  $X_{j-1}C_{j-1}, X_{j-1}D_{j-1}$ , and its last edge is one of  $A_jX_j, B_jX_j$ . As noted above, the turn formed by the last edge of  $\gamma_j$  and the first edge of  $\gamma_{j+1}$  is evil. We conclude that the loop  $\gamma_1\gamma_2\cdots\gamma_n$  is evil, contradicting rigidity.  $\square$


The preceding result shows that there is a well-defined order relation on the set of skeletons in the support of a rigid measure  $m$  defined as follows:  $S \prec S'$  if there exist skeletons  $S_1, S_2, \dots, S_k$  such that

$$S = S_1 \prec_0 S_2 \prec_0 \cdots \prec_0 S_k = S'.$$

The following figure shows the support of a rigid measure  $m \in \mathcal{M}_6$ . The elements of a maximal collection of mutually inequivalent root edges have been indicated with dots.



For this measure, there is a smallest skeleton (relative to  $\prec$ ) pictured below. The reader can easily draw all the other skeletons and determine the order relation.



#### 4. HORN INEQUALITIES AND CLOCKWISE OVERLAYS

The results of [21, 22] show that the triples  $(\alpha, \beta, \gamma) = \partial m$  with  $m \in \mathcal{M}_r$  are precisely those triples of increasing nonnegative vectors in  $\mathbb{R}^r$  with the property that there exist selfadjoint matrices  $A, B, C \in M_r(\mathbb{C})$  such that

$$\lambda_A(\ell) = \alpha_{r+1-\ell}, \lambda_B(\ell) = \beta_{r+1-\ell}, \lambda_C(\ell) = \gamma_{r+1-\ell}, \quad \ell = 1, 2, \dots, r,$$

and  $A+B+C$  is a multiple of the identity, namely,  $2\omega(m)1_{\mathbb{C}^r}$ . The Horn inequalities for these matrices are

$$\sum_{i \in I} \lambda_A(i) + \sum_{j \in J} \lambda_B(j) + \sum_{k \in K} \lambda_C(k) \leq 2s\omega(m)$$

when  $I, J, K \subset \{1, 2, \dots, r\}$  have cardinality  $s$  and  $c_{IJK} > 0$ . Applying this inequality to the matrices  $-A, -B, -C$  instead and switching signs, we obtain

$$\sum_{i \in I} \lambda_A(r+1-i) + \sum_{j \in J} \lambda_B(r+1-j) + \sum_{k \in K} \lambda_C(r+1-k) \geq 2s\omega(m)$$

Equivalently,

$$\sum_{i \in I} \alpha_i + \sum_{j \in J} \beta_j + \sum_{k \in K} \gamma_k \geq 2s\omega(m).$$

Now, the sets  $I, J, K$  are obtained from some measure  $\nu \in \mathcal{M}_s$  with weight  $r-s$ , and we will see how this inequality follows from the superposition of the support of  $m$  and the puzzle associated with  $\nu$ . Let  $h$  be the honeycomb provided by Lemma 1.1. Let  $D \subset \Delta_r$  be a region bounded by small edges, and let  $X_j Y_j, j = 1, 2, \dots, p$  be an enumeration of the edges of  $\partial D$ , oriented so that  $D$  lies on the left of  $X_j Y_j$ . For each  $j$ , there is  $\varepsilon_j = \varepsilon_{X_j Y_j} = \pm 1$  such that  $Y_j - X_j \in \{\varepsilon_j u, \varepsilon_j v, \varepsilon_j w\}$ . The definition of honeycombs implies then the identity

$$\sum_{XY \subset \partial D} \varepsilon_{XY} h(XY) = \sum_{j=1}^p \varepsilon_j h(X_j Y_j) = 0,$$

which is easily deduced by induction on the size of  $D$ . We would like to verify that the sum

$$S = \sum_{i \in I} h(A_{i-1} A_i) + \sum_{j \in J} h(B_{j-1} B_j) + \sum_{k \in K} h(C_{k-1} C_k)$$

is nonnegative, where  $I, J, K$  are given by a measure  $\nu \in \mathcal{M}_s$  with  $\omega(\nu) = r-s$ . The inflation of the measure  $\nu$  yields a partition of  $\Delta_r$  into white pieces (the translated parts of  $\Delta_s$ ), gray parallelograms, and light gray pieces. Denote by  $D$  the union of the gray parallelograms  $P_1, P_2, \dots, P_\sigma$  and white pieces  $W_1, W_2, \dots, W_\tau$ . The edges  $A_{i-1} A_i, i \in I, B_{j-1} B_j, j \in J$ , and  $C_{k-1} C_k, k \in K$  are contained in the intersection  $\partial D \cap \partial \Delta$ . Since

$$\sum_{XY \subset \partial D} \varepsilon_{XY} h(XY) = \sum_{XY \subset \partial W_\ell} \varepsilon_{XY} h(XY) = 0, \quad \ell = 1, 2, \dots, \tau,$$

and

$$S = \sum_{\text{white } XY \subset \partial D \cap \partial \Delta_r} \varepsilon_{XY} h(XY),$$

we have


$$\begin{aligned} S &= S - \sum_{XY \subset \partial D} \varepsilon_{XY} h(XY) + \sum_{\ell=1}^{\tau} \sum_{XY \subset \partial W_\ell} \varepsilon_{XY} h(XY) \\ &= - \sum_{\ell=1}^{\sigma} \sum_{\text{gray } XY \subset \partial P_\ell} \varepsilon_{XY} h(XY), \end{aligned}$$

where the last sums are taken over the light gray edges of  $P_\ell$ . Condition (3) implies that


$$- \sum_{\text{gray } XY \subset \partial P_\ell} \varepsilon_{XY} h(XY) = \sum_e m(e),$$

where the sum is taken over all small edges contained in  $P_\ell$  which are not parallel to the sides of  $P_\ell$ . This immediately implies the desired Horn inequality, and it also tells us when equality is attained: this happens if and only if all the edges  $e \subset P_\ell$  for which  $m(e) > 0$  are parallel to the edges of  $P_\ell, \ell = 1, 2, \dots, s$ . In other words,

the support of  $m$  must cross each  $P_\ell$  along lines parallel to the edges of  $P_\ell$ . The following figure illustrates the support of a measure  $\nu$ , the inflation of  $\nu$ , and the support of a measure  $m$  which satisfies the Horn equality associated with  $\nu$ . In this example the support of the measure  $m$  never crosses the white puzzle pieces.



The next example involves basically the mirror image of the measure  $m$ , and its support never crosses the light gray pieces.



This phenomenon is related with the fact that the support of  $m$  is actually a rigid skeleton in both cases. A rigid skeleton does not cross itself transversely. Thus, in a case of equality, it cannot have any branch points in the interior of a gray parallelogram. It follows then that the skeleton always crosses these parallelograms between white pieces or between gray pieces, but not both.

Assume that we are in a case of equality


$$\sum_{i \in I} h(A_{i-1}A_i) + \sum_{j \in J} h(B_{j-1}B_j) + \sum_{k \in K} h(C_{k-1}C_k) = 0.$$

In this case, we can define a measure  $\mu \in \mathcal{M}_s$  by moving the support of  $m$  to  $\Delta_s$  in the following way: those parts which are contained in white puzzle pieces are simply translated back to  $\Delta_s$  (along with their densities); the segments in the support of  $m$  which cross between white pieces are deleted; the segments which cross between light gray pieces are replaced by the corresponding parallel sides of white pieces, and the density is preserved. It may be that several segments cross between light gray pieces, in which case the density of the corresponding side of a white piece is the sum of their densities. When the measure  $\mu$  can be obtained using this procedure, we will say that  $\mu$  is obtained by *contracting*  $m$ , and that  $\mu$  is *clockwise* from  $\nu$  (or that  $(\mu, \nu)$  form a *clockwise overlay*). This is easily seen to be an extension of the notion of clockwise overlay introduced in [22] (see also item (1), second case, in the proof of Theorem 4.2). Generally, a clockwise overlay  $(\mu, \nu)$  can be obtained



by shrinking more than one measure  $m$ . Indeed, the shrinking operation loses all the information about the branch points of  $m$  in the light gray puzzle pieces.

In the first case illustrated above, the support of the measure  $\mu$  is actually contained in the support of  $\nu$ ; this is what happens when the support of  $m$  does not cross the white pieces. In the second case illustrated above we obtain the following figure for the supports of  $\nu$  and  $\mu$ .





We will need one more important property of clockwise overlays.

**Proposition 4.1.** *Let  $(\mu, \nu)$  be a clockwise overlay obtained by contracting a measure  $m$ . Then  $\omega(\mu) = \omega(m)$ , and  $\mu^* \leq m^*$ .*

*Proof.* Consider the puzzle obtained by inflating the measure  $\varepsilon\nu$  for  $\varepsilon > 0$ . The white pieces of the puzzle are independent of  $\varepsilon$ . Since the support of  $m$  intersects any gray parallelogram in the puzzle of  $\nu$  only on segments parallel to the edges of the parallelogram, it follows that there exists a measure  $m_\varepsilon$  obtained by translating the support of  $m$  in each white piece, and applying appropriate translation and/or shrinking in the gray parallelograms and light gray puzzle pieces. Clearly  $m_1 = m$ , and all the measures  $m_\varepsilon$  are homologous to  $m$ ; in fact, homologous sides have equal densities, and therefore  $\omega(m) = \omega(m_\varepsilon)$  for all  $\varepsilon > 0$ . (Here it may be useful to recall that  $\omega(m)$  is defined in terms of its densities outside  $\Delta_r$ , and the outside edges are not generally present in our drawings.) Moreover, all the measures  $m_\varepsilon^*$  have the same support, except that some of the densities are decreased for  $\varepsilon < 1$ . The measure  $\mu$  is simply the limit of  $m_\varepsilon$  as  $\varepsilon \rightarrow 0$ , and the statement follows immediately from this observation.

The following pictures illustrates the process as applied to the above examples for  $\varepsilon = 2/3$  and  $\varepsilon = 1/3$ .





□

Unfortunately, the definition of clockwise overlays is not quite explicit since they are seen as the result of a process — something akin to defining a car as the end product of car manufacture. We can however use the relation  $\prec_0$  between skeletons to produce an important class of clockwise overlays.


**Theorem 4.2.** *Let  $\mu_1, \mu_2 \in \mathcal{M}_r$  be such that  $\mu_1 + \mu_2$  is rigid, the support  $S_j$  of  $\mu_j$  is a skeleton, and  $S_2 \not\prec_0 S_1$ . Then  $(\mu_1, \mu_2)$  is a clockwise overlay.*

*Proof.* We need to inflate  $\mu_2$ , and construct a measure  $m_1$  such that  $\mu_1$  is obtained from  $m_1$  by the shrinking process described above. It is clear what the measure  $m_1$  should be on the interior of every white puzzle piece. The common edges of  $S_1$  and  $S_2$  cannot be root edges; orient them away from the root edges, and attach them (along with their  $\mu_1$  masses) to the white puzzle piece on their right side. What remains to be proved is that this partially defined measure can be extended so as to satisfy the balance condition at all points. For this purpose we only need to analyze the situation at lattice points where  $S_1$  and  $S_2$  meet. For each such lattice point, there will be 2, 3, or 4 edges of each skeleton meeting at that point, and this gives rise to many possibilities. In order to reduce the number of cases we need to study, observe that the inflation construction is invariant relative to rotations of  $60^\circ$ , and therefore the position (but perhaps not the orientation) of the edges in one of the skeletons can be fixed. In the following figures, the arrows indicate the orientation on the edges in  $S_1 \cap S_2$ . The other edges of  $S_1$  are dashed, and the other edges of  $S_2$  are solid without arrows. In each case, the extension required after inflation is indicated by dashed lines crossing (or on the boundary of) parallelogram pieces. In the following enumeration, the label  $(p, q)$  signifies that  $S_1$  has  $p$  and  $S_2$  has  $q$  edges meeting at one point.

- (1) (2, 2) The edges of the skeletons may overlap, and after a rotation the orientation is as in the figure below.



No extensions are required in this case. If the skeletons do not overlap, we have two possibilities:




and finally




which would imply  $S_2 \prec_0 S_1$ , contrary to the hypothesis.


- (2) (3,2) In this case there is (up to rotations) only the case illustrated in the figure.




- (3) (4,2) Up to rotations, there are three possibilities. In the first one we have an extension as shown.



The orientation shown above is the only one which is compatible with the rigidity of  $m$ . In the following figure, the orientation given is also the only possible one.




The third situation




implies  $S_2 \prec S_1$ .


- (4) (2,3) The case illustrated is the only one up to rotations.




- (5) (3,3) There are two cases up to rotations.




The second case is not compatible with rigidity.




- (6) (4, 3) There is only one position of  $S_1$  compatible with rigidity, but there are two possible orientations.




The second orientation requires a different extension.




- (7) (2, 4) There are three possibilities up to rotation.




In the figure above, there is no ambiguity in the orientation. For the illustration we assigned  $\mu_2$  masses of 1 and 2 to the edges.




The orientation is also clear in this case. The third case implies  $S_2 \prec S_1$ .




- (8) (3, 4) There is only one position compatible with rigidity, and there are only two possible orientations.






- (9) (4,4) When the two skeletons overlap completely, there are two possible orientations.



When the skeletons do not overlap completely, there is only one relative position of the two skeletons which is compatible both with rigidity and with  $S_2 \not\prec S_1$ . There is only one possible orientation.



□

The following result follows easily by induction, inflating successively the measures  $\mu_p, \mu_{p-1}, \dots, \mu_{t+1}$ .

**Corollary 4.3.** *Let  $m \in \mathcal{M}_r$  be a rigid measure, and write it as  $m = \sum_{\ell=1}^p \mu_\ell$ , where  $\mu_\ell$  is supported on the skeleton  $S_\ell$ . Assume also that  $S_i \prec S_j$  implies that  $i \leq j$ . Then the pair  $(\sum_{\ell=1}^t \mu_\ell, \sum_{\ell=t+1}^p \mu_\ell)$  is a clockwise overlay for  $1 \leq t < p$ .*

For the clockwise overlays  $(\mu_1, \mu_2)$  considered in the preceding two results there is a canonical construction for the measure  $m_1 \in \mathcal{M}_{r+\omega(\mu_2)}$ . We will call this measure  $m_1$  the *stretch of  $\mu_1$  to the puzzle of  $\mu_2$* .

## 5. PROOF OF THE MAIN RESULTS

Fix a triple  $(I, J, K)$  of subsets with cardinality  $r$  of  $\{0, 1, \dots, n\}$  such that  $c_{IJK} = 1$ , and let  $m \in \mathcal{M}_r$  be the corresponding measure. It will be convenient now to use the normalization  $\tau(1) = n$  in a finite factor. This will not require

renormalizations when passing to a subfactor, and has the added benefit of working in finite dimensions as well. In order to prove the intersection results in the introduction, we will want to prove the following related properties:

**Property  $A(I, J, K)$  or  $A(m)$ .** Given a  $\Pi_1$  factor  $\mathcal{A}$  with  $\tau(1) = n$ , and given flags  $\mathcal{E}, \mathcal{F}, \mathcal{G}$  with  $\tau(E_\ell) = \tau(F_\ell) = \tau(G_\ell) = \ell$ ,  $\ell = 0, 1, 2, \dots, n$ , the intersection

$$S(\mathcal{E}, I) \cap S(\mathcal{F}, J) \cap S(\mathcal{G}, K)$$

is not empty.

**Property  $B(I, J, K)$  or  $B(m)$ .** There exists a lattice polynomial  $p \in L(\{e_j, f_j, g_j : 1 \leq j \leq n\})$  with the following property: for any finite factor  $\mathcal{A}$  with  $\tau(1) = n$ , and for generic flags  $\mathcal{E} = (E_j)_{j=0}^n$ ,  $\mathcal{F} = (F_j)_{j=0}^n$ ,  $\mathcal{G} = (G_j)_{j=0}^n$  such that  $\tau(E_j) = \tau(F_j) = \tau(G_j) = j$ , the projection  $P = p(\mathcal{E}, \mathcal{F}, \mathcal{G})$  has trace  $\tau(P) = r$  and, in addition

$$\tau(P \wedge E_i) = \tau(P \wedge F_j) = \tau(P \wedge G_k) = \ell$$

when  $i_\ell \leq i < i_{\ell+1}, j_\ell \leq j < j_{\ell+1}, k_\ell \leq k < k_{\ell+1}$  and  $\ell = 0, 1, \dots, r$ , where  $i_0 = j_0 = k_0 = 0$  and  $i_{r+1} = j_{r+1} = k_{r+1} = n+1$ .

We will prove these properties by reducing them to simpler measures for which they are trivial. The basic reduction is from an arbitrary measure to a skeleton.

**Proposition 5.1.** *Let  $m \in \mathcal{M}_r$  a rigid measure, and write  $m = \sum_{\ell=1}^p \mu_\ell$ , where  $\mu_\ell$  is supported by a skeleton  $S_\ell$ , and  $S_i \prec S_j$  implies  $i \leq j$ . Let  $\tilde{\mu}_1 \in \mathcal{M}_{\tilde{r}}$ ,  $\tilde{r} = \sum_{\ell=2}^p \omega(\mu_\ell)$  be the stretch of  $\mu_1$  to the puzzle of  $m' = \sum_{\ell=2}^p \mu_\ell$ . If  $A(\tilde{\mu}_1)$  and  $A(m')$  (resp.,  $B(\tilde{\mu}_1)$  and  $B(m')$ ) are true, then  $A(m)$  (resp.,  $B(m)$ ) is true as well.*

*Proof.* With the usual notation  $A_i = iu$ ,  $X_i = A_i + w$ , the edges  $A_i X_i$  are oriented in the direction of  $w$  (if they belong to the support of  $m$ ). Let us set  $a_i = m(A_i X_i)$ ,  $a_i^{(1)} = \mu_1(A_i X_i)$ , and  $a'_i = m'(A_i X_i)$ , so that  $a_i = a'_i + a_i^{(1)}$ , and  $n = r + \sum_{i=0}^r a_i$ . The measure  $\tilde{\mu}_1$  is associated with the triangle  $\Delta_{r_1}$ , where  $r_1 = r + \sum_{i=0}^r a'_i$ . Its support may intersect the left side of this triangle only at the points  $A_{\ell(i)}$ , where  $\ell(0) = 0$ , and  $\ell(i) = i + \sum_{s=0}^{i-1} a'_s$  for  $i > 0$ ; this follows from the way the inflation of  $m'$  is constructed, and from the outward orientation of the segments  $A_i X_i$ . Moreover,  $\tilde{\mu}_1(A_{\ell(i)} X_{\ell(i)}) = a_i^{(1)}$  for  $i = 0, 1, \dots, r$ . Denote by  $I^{(1)}, J^{(1)}, K^{(1)} \subset \{1, 2, \dots, n\}$  the sets determined by the measure  $\tilde{\mu}_1$ , and by  $I', J', K' \subset \{1, 2, \dots, n_1\}$  those corresponding with  $m'$ , where we set  $n_1 = n - \omega(m_1)$ . For instance, we have

$$I^{(1)} = \bigcup_{j=1}^r \left\{ s + \sum_{\ell=0}^{j-1} a_\ell^{(1)} + \sum_{\ell=0}^{j-2} (a'_\ell + 1) : s = 1, 2, \dots, a'_{j-1} + 1 \right\},$$

where the second sum is zero for  $j = 1$ , and

$$I' = \{i_t - \sum_{\ell=0}^{t-1} a_\ell^{(1)} : t = 1, 2, \dots, r\},$$

where  $I = \{i_1, i_2, \dots, i_r\}$ . Observe that  $i_\ell^{(1)} = i_t$  for  $\ell = t + \sum_{s=0}^{t-1} a'_s = i'_t$ .

Assume first that  $A(\tilde{\mu}_1)$  and  $A(m')$  are true, and let  $\mathcal{E}, \mathcal{F}, \mathcal{G}$  be arbitrary flags in a  $\Pi_1$  factor such that  $\tau(E_i) = \tau(F_i) = \tau(G_i) = i$  for  $i = 0, 1, \dots, n$ , and  $\tau(1) = n$ . Property  $A(\tilde{\mu}_1)$  implies the existence of a projection  $P_1 \in \mathcal{A}$  such that  $\tau(P_1) = r_1$  and

$$\tau(P_1 \wedge E_{i_\ell}) \geq \ell, \ell = 1, 2, \dots, r_1.$$

As noted above, we have  $i_\ell^{(1)} = i_t$  for  $\ell = i'_t$ , and therefore we have

$$\tau(P_1 \wedge E_{i'_t}) \geq t + \sum_{s=0}^{t-1} a'_s = i'_t, \quad t = 1, 2, \dots, r,$$

with analogous inequalities for  $\mathcal{F}$  and  $\mathcal{G}$ . Consider now the factor  $\mathcal{A}_1 = P_1 \mathcal{A} P_1$  with the trace  $\tau_1 = \tau|_{\mathcal{A}_1}$ , so that  $\tau_1(1_{\mathcal{A}'_1}) = \tau(P_1) = r_1$ . The inequalities above imply the existence of a flag  $\mathcal{E}'$  in  $\mathcal{A}_1$  such that  $\tau_1(E'_j) = j$  for  $j = 1, 2, \dots, n_1$ , and

$$E'_{i'_p} \leq P_1 \wedge E_{i_p}, \quad p = 1, 2, \dots, r.$$

Analogous considerations lead to the construction of flags  $\mathcal{F}'$  and  $\mathcal{G}'$ . Property  $A(m')$  implies now the existence of a projection  $P \in \mathcal{A}_1$  such that  $\tau_1(P) = r$ ,

$$\tau_1(P \wedge E'_{i'_p}) \geq p, \quad p = 1, 2, \dots, r,$$

and analogous inequalities are satisfied for  $\mathcal{F}'$  and  $\mathcal{G}'$ . Clearly the projection  $P$  satisfies

$$\tau(P \wedge E_{i_p}) \geq p, \quad p = 1, 2, \dots, r,$$

so that it solves the intersection problem for the sets  $I, J, K$ .

The case of property  $B$  is settled analogously. The difference is that  $P_1$  is given as a lattice polynomial  $P_1 = p_1(\mathcal{E}, \mathcal{F}, \mathcal{G})$ , and the projections  $E'_j$  can be taken to be of the form  $P_1 \wedge E_i$ , and hence they too are lattice polynomials in  $\mathcal{E}, \mathcal{F}, \mathcal{G}$ . Finally, the solution  $P$  is given as  $P = p'(\mathcal{E}', \mathcal{F}', \mathcal{G}')$ , where the existence of  $p'$  is given by property  $B(m')$ . One must however assume that  $\mathcal{E}', \mathcal{F}', \mathcal{G}'$  are generic flags, and this simply amounts to an additional genericity condition on the original flags.  $\square$

The preceding proposition shows that proving property  $A(m)$  or  $B(m)$  can be reduced to proving it for simpler measures, at least when  $m$  is not extremal. A dual reduction is obtained by recalling that a projection  $P$  belongs to  $S(\mathcal{E}, I)$  if and only  $P^\perp = 1 - P$  belongs to  $S(\mathcal{E}^\perp, I^*)$ . Moreover, if the sets  $I, J, K$  are associated to the measure  $m \in \mathcal{M}_r$ , then  $I^*, J^*, K^*$  are the sets associated to the measure  $m^*$ . Therefore  $A(m)$  is equivalent to  $A(m^*)$  and  $B(m)$  is equivalent to  $B(m^*)$ .

To quantify these reductions, we define for each measure  $m \in \mathcal{M}_r$  the positive integer  $\kappa(m)$  as the number of gray parallelograms in the puzzle obtained by inflating  $m$ . This is equal to the number of white piece edges which have positive measure. Analogously, for  $m \in \mathcal{M}_r^*$ , we define  $\kappa^*(m)$  to be the number of gray parallelograms in the puzzle obtained by \*inflating  $m$ . With this definition it is clear that

$$\kappa(m) = \kappa^*(m^*), \quad m \in \mathcal{M}_r.$$

Indeed, the two numbers count pieces of the same puzzle.

With the notation of the preceding proposition, we have

$$\kappa(\widetilde{\mu}_1) = \kappa(\mu_1) < \kappa(m), \quad \kappa(m') < \kappa(m),$$

unless  $m = \mu_1$ . Indeed,  $\kappa(\widetilde{\mu}_1) = \kappa(\mu_1)$  because  $\mu_1$  and  $\widetilde{\mu}_1$  are homologous, and the supports of  $\mu_1$  and  $m'$  are strictly contained in the support of  $m$ . In fact, the support of  $m'$  does not contain the root edges of  $\mu_1$ , and the support of  $\mu_1$  does not contain the root edges of the extremal summands of  $m'$ . Thus the preceding proposition also allows us to reduce the proof of these properties to measures with smaller values of  $\kappa$  in case either  $m$  or  $m^*$  is not extremal. The exceptional situations

in which both  $m$  and  $m^*$  are extremal are very few in number. To see this we need to use the structure of the convex polyhedral cone

$$C_r = \{\partial m : m \in \mathcal{M}_r\},$$

whose facets were determined in [22]. If  $\partial m = (\alpha, \beta, \gamma)$ , these facets are of two kinds. The first kind are the *chamber facets* determined by an equality of the form  $\alpha_\ell = \alpha_{\ell+1}$ ,  $\beta_\ell = \beta_{\ell+1}$ ,  $\gamma_\ell = \gamma_{\ell+1}$  for  $1 \leq \ell < r$  or  $\alpha_r = \omega(m)$ ,  $\beta_r = \omega(m)$ ,  $\gamma_r = \omega(m)$ . The second kind are the *regular facets* determined by Horn identities

$$\sum_{i \in I} \alpha_i + \sum_{j \in J} \beta_j + \sum_{k \in K} \gamma_k = \omega(m),$$

where  $I, J, K \subset \{1, 2, \dots, r\}$  have  $s < r$  elements and  $c_{IJK} = 1$ .

For a given measure  $m \in \mathcal{M}_r$ , we define the number of attachment points  $\Gamma(m)$  to be the number of chamber facets to which  $m$  does not belong. The reason for this terminology is that  $\Gamma(m)$  is precisely the number of points on the sides of  $\Delta_r$  which are endpoints of interior edges in the support of  $m$ . The vertices of  $\Delta_r$  should also counted as attachment points when they are branch points of the measure.


**Proposition 5.2.** *Let  $m \in \mathcal{M}_r$  be an extremal rigid measure. If  $m^*$  is extremal as well, then  $\Gamma(m) = 1$ .*

*Proof.* Assume that  $m$  and  $m^*$  are both extremal, and  $\Gamma(m) > 1$ . Note first that  $\partial m$  is extremal in  $C_r$ . Indeed, in the contrary case, we would have  $\partial m = \partial m_1 + \partial m_2$  with  $\partial m_1$  not a positive multiple of  $\partial m$ . This would however imply  $m = m_1 + m_2$  by rigidity, and hence  $m_1$  is a multiple of  $m$ , a contradiction.

Next, since  $\Gamma(m) = \Gamma(m')$  for homologous  $m, m'$ , we may assume that  $m^* = \mu_e$  for some root edge  $e$ . Indeed,  $m^* = m^*(e)\mu_e$  is homologous to  $\mu_e$ , and therefore  $m$  is homologous to  $\mu_e^*$ .

The definition of  $\Gamma(m)$  implies that  $\partial m$  belongs to  $3r - \Gamma(m) = \dim C_r - \Gamma(m)$  chamber facets. However, an extremal measure must belong to at least  $\dim C_r - 1$  facets, and hence  $\partial m$  belongs to at least one regular facet. As seen earlier, there must then exist a clockwise overlay  $(m_1, m_2)$  such that  $m_1$  is obtained by contracting  $m$ . It follows from Proposition 4.1 that  $0 \neq m_1^* \leq m^*$ . Since  $m_1^*$  has integer densities, we must have  $m_1^* = m^*$ , and this implies that  $m_1 = m$ , a contradiction.  $\square$

Thus the repeated application of the reduction procedure to  $m$  and  $m^*$  leads eventually to one of the three skeletons pictured below.



For these, the intersection problem is completely trivial. Indeed, consider the first of the three on  $\Delta_r$ , and with  $\omega(m) = s$ . We have then  $I = \{1, 2, \dots, r\}$  and  $J = K = \{s+1, s+2, \dots, s+r\}$ , and the desired element in

$$S(\mathcal{E}, I) \cap S(\mathcal{F}, J) \cap S(\mathcal{G}, K)$$



is simply  $E_r$ . Thus  $A(m)$  is true for this measure. To show that  $B(m)$  is true as well, we must verify that generically we also have

$$\tau(E_r \wedge F_\ell) = \tau(E_r \wedge G_\ell) = \max\{0, r + \ell - n\}$$

for  $\ell = 1, 2, \dots, n$ . This follows easily from the following result.

**Proposition 5.3.** *Let  $E$  and  $F$  be two projections in a finite factor  $\mathcal{A}$ . There is an open dense set  $O \subset \mathcal{U}(\mathcal{A})$  such that*

$$\tau(E \wedge UFU^*) = \max\{0, \tau(E) + \tau(F) - \tau(1)\}$$

for  $U \in O$ .

*Proof.* Replacing  $E$  and  $F$  by  $E^\perp$  and  $F^\perp$  if necessary, we may assume that  $\tau(E) + \tau(F) \leq \tau(1)$ . Since  $\mathcal{A}$  is a factor, we can replace  $F$  with any other projection with the same trace. In particular, we may assume that  $F \leq E^\perp$ . The condition  $\tau(E \wedge UFU^*) = 0$  is satisfied if the operator  $FUF$  is invertible on the range of  $F$ . The proposition follows because the set  $O$  of unitaries satisfying this condition is a dense open set in  $\mathcal{U}(\mathcal{A})$ . To verify this fact, it suffices to consider the case in which the algebra  $\mathcal{A}$  is of the form  $\mathcal{A} = \mathcal{B} \otimes M_2(\mathbb{C})$  for some other finite factor  $\mathcal{B}$ , and

$$F = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$$

An arbitrary unitary  $U \in \mathcal{A}$  can be written as

$$U = \begin{bmatrix} T & (I - TT^*)^{1/2}W \\ V(I - T^*T)^{1/2} & -VT^*W \end{bmatrix},$$

where  $T, V, W \in \mathcal{B}$ ,  $V$  and  $W$  are unitary, and  $\|T\| \leq 1$ . Since  $\mathcal{B}$  is a finite von Neumann algebra,  $T$  can be approximated arbitrarily well in norm by an invertible operator  $T'$ , in which case  $U$  is approximated in norm by the operator

$$U' = \begin{bmatrix} T' & (I - T'T'^*)^{1/2}W \\ V(I - T'^*T')^{1/2} & -VT'^*W \end{bmatrix}$$

with  $FU'F$  invertible. In finite dimensions, the complement of  $O$  is defined by the single homogeneous polynomial equation  $\det(FUF + F^\perp) = 0$ . Thus  $O$  is open in the Zariski topology.  $\square$

**Corollary 5.4.** *Properties  $A(m)$  and  $B(m)$  are true for all rigid measures  $m$ .*

This proves finally Theorems 0.4 and 0.5. The fact that Theorem 0.3 follows from Theorem 0.4 was already shown in [4].


## 6. SOME ILLUSTRATIONS

We have just seen that proving property  $A(I, J, K)$  or  $B(I, J, K)$  can be reduced, in case  $c_{IJK} = 1$ , to the case in which the associated measure  $m$  has precisely one attachment point. We will illustrate how this reduction works in a few cases.

Given a measure  $m \in \mathcal{M}_r$ , a point  $A_\ell$ ,  $\ell = 1, 2, \dots, r$ , is an attachment point of  $m$  precisely when  $m(A_\ell X_\ell) > 0$ . The solution to the associated Schubert intersection problem will only depend on the projections  $E_{i(\ell)}$  where  $\ell$  is an attachment point. These projections, and the analogous  $F_{j(\ell)}, G_{k(\ell)}$ , will be called the *attachment projections* for the problem. With the notation Proposition 5.1, the attachment projections of  $\widetilde{\mu}_1$  are exactly the same as those of  $\mu_1$ , and are therefore among the attachment projections of  $m$ . The attachment projections of  $m^*$  are of the


form  $P^\perp = 1 - P$ , where  $P$  is an attachment projection for  $m$ . These observations allow us to construct solutions to intersection problems without actually having to construct the measure  $\widetilde{\mu}_1$  and focus instead on the attachment projections of  $\mu_1$ .

We proceed now to solve the intersection problems associated with some skeletons. Consider first an extreme measure  $m \in \mathcal{M}_r$  with two attachment points. The following picture shows the supports of  $m$  and  $m^*$ .



For the illustration we took  $r = 3$  and density 3 on the support, but the results will hold for the general case. Note that  $m^*$  is a sum of two extremal measures with one attachment point each. If  $X$  and  $Z$  are the attachment projections of  $m$ , the attachment projections of these skeletons are  $X^\perp$  and  $Z^\perp$ . Neither of the two skeletons precedes the other, and following the method of Proposition 5.1, we see that the solution of the intersection problem associated with  $m^*$  is generically  $X^\perp \wedge Z^\perp$ . It follows that the intersection problem associated with  $m$  has the generic solution  $X \vee Y$ .


There are two kinds of skeletons with three attachment points. The first one, and its dual, are illustrated below.



Assume that the attachment projections are  $X, Y$  and  $Z$ . As in the preceding situation,  $m^*$  is a sum of three extremal measures with one attachment point, and there are no precedence relations among the skeletons. It follows that the generic solution of the intersection problem is  $X \vee Y \vee Z$ .

The two cases just mentioned correspond to the reductions considered in [25] for finite dimensions, and in [9] for the factor case. Note however that these papers also apply these reductions when  $c_{IJK} > 1$ .

Consider next the other kind of skeleton with three attachment points, and with attachment projections  $X, Y, Z$ .



In this case,  $m^*$  is the sum of three extremal measures with two attachment points each, and with no precedence relations. The intersection problems associated with the three skeletons have then generic solutions  $X^\perp \vee Y^\perp$ ,  $X^\perp \vee Z^\perp$ , and  $Y^\perp \vee Z^\perp$ . According to Proposition 5.1, the solution of the intersection problem for  $m^*$  will be (generically) the intersection of these three projections, so that the problem


associated with  $m$  has the solution

$$(X \wedge Y) \vee (X \wedge Z) \vee (Y \wedge Z).$$

Several of the proofs of Horn inequalities in the literature can now be deduced by considering rigid measures which are sums of extremal measures with 1, 2 or 3 attachment points. Consider, for instance, a measure  $m \in \mathcal{M}_r$  defined by

$$m = \rho + \sum_{\ell=1}^r (\mu_\ell + \nu_\ell),$$

where  $\rho$  has attachment point  $C_r$ ,  $\mu_1$  has attachment point  $A_r$ ,  $\nu_1$  has attachment point  $B_r$ ,  $\mu_\ell$  has attachment points  $A_{r-\ell+1}$  and  $C_{\ell-1}$ , and  $\nu_\ell$  has attachment points  $B_{r-\ell+1}$  and  $C_{\ell-1}$  for  $\ell > 1$ .



The only precedence relations are  $\mu_\ell \prec_0 \nu_k$  and  $\nu_\ell \prec_0 \mu_k$  for  $\ell < k$ . Generically, the associated intersection problem is solved as follows. Set  $P_0 = G_r$  and

$$P_{\ell+1} = [(G_\ell \wedge P_\ell) \vee (F_{r-\ell} \wedge P_\ell)] \wedge [(G_\ell \wedge P_\ell) \vee (E_{r-\ell} \wedge P_\ell)]$$

for  $\ell = 1, 2, \dots, r-1$ . The space  $P_r$  is the generic solution. The sets  $I, J, K$  associated with  $m$  are easily calculated. Using the notations

$$c = \omega(\rho), \quad a_\ell = \omega(\mu_\ell), \quad b_\ell = \omega(\nu_\ell) \quad \text{for } \ell = 1, 2, \dots, r,$$

we have

$$n = r + c + \sum_{\ell=1}^r (a_\ell + b_\ell),$$

and  $I = \{n + 1 - (a_1 + a_2 + \dots + a_\ell + \ell) : \ell = 1, 2, \dots, r\}$ ,  $J = \{n + 1 - (b_1 + b_2 + \dots + b_\ell + \ell) : \ell = 1, 2, \dots, r\}$ , and  $K = \{a_1 + b_1 + \dots + a_\ell + b_\ell + \ell : \ell = 1, 2, \dots, r\}$ . These sets yield the eigenvalue inequalities proved in [23].


Consider next sequences of integers

$$0 \leq z_1 \leq z_2 \leq \dots \leq z_p, \quad 0 \leq w_1 \leq w_2 \leq \dots \leq w_p$$

such that  $z_p + w_p \leq r$ , and consider the measure  $m \in \mathcal{M}_r$  defined by

$$m = \sum_{i=1}^p \mu_i,$$

where  $\mu_\ell$  has attachment points  $A_{z_\ell}, B_{w_\ell}$ , and  $C_{r-z_\ell-w_\ell}$ .



The illustration uses  $p = 3$ ,  $r = 6$ ,  $z_1 = 1, z_2 = 2, z_3 = 3$ ,  $w_1 = w_2 = 1$ , and  $w_3 = 2$ . We have  $\mu_\ell \prec_0 \mu_k$  only when  $\ell < k$ ,  $w_\ell < w_k$  and  $z_\ell < z_k$ . If we set  $P_1 = E_{z_1} \vee F_{w_1} \vee G_{r-z_1-w_1}$  and

$$P_{\ell+1} = (E_{z_{\ell+1}} \wedge P_\ell) \vee (F_{w_{\ell+1}} \wedge P_\ell) \vee (G_{r-z_{\ell+1}-w_{\ell+1}} \wedge P_\ell)$$

for  $\ell = 1, 2, \dots, d-1$ , then  $P_d$  is the generic solution of the intersection problem. Assume that  $\omega(\mu_i) = 1$  for all  $i$ , and use the notation

$$1_{x < y} = \begin{cases} 1 & \text{if } x < y, \\ 0 & \text{if } x \geq y. \end{cases}$$

Then for the corresponding intersection problem we have  $n = r + p$ ,  $I(\ell) = \ell + \sum_{i=1}^p 1_{z_i < \ell}$ ,  $J(\ell) = \ell + \sum_{i=1}^p 1_{w_i < \ell}$ , and  $n + 1 - K(r + 1 - \ell) = \ell + \sum_{i=1}^p 1_{w_i + z_i < \ell}$  for  $\ell = 1, 2, \dots, r$ . These sets yield the eigenvalue inequalities proved in [24].


One can produce such families of inequalities using more complicated skeletons. Observe for instance that, given integers  $a, b, c, d$  such that  $a + b + c + d = r$ , there exists a skeleton in  $\Delta_r$  with attachment points  $A_a, A_{a+b+c}, B_{b+d}$ , and  $C_{c+d}$ . Call  $\mu_{a,b,c,d}$  the smallest extremal measure with integer densities supported by this skeleton. A measure of the form

$$m = \sum_{\ell=1}^p \mu_{a_\ell, b_\ell, c_\ell, d_\ell}$$


will be rigid if the following conditions are satisfied:

$$a_\ell \leq a_{\ell+1}, d_\ell \leq d_{\ell+1}, c_\ell + d_\ell \leq c_{\ell+1} + d_{\ell+1}, b_\ell + d_\ell \leq b_{\ell+1} + d_{\ell+1}$$


for  $\ell = 1, 2, \dots, p-1$ . Moreover,  $\mu_{a_\ell, b_\ell, c_\ell, d_\ell} \prec \mu_{a_{\ell'}, b_{\ell'}, c_{\ell'}, d_{\ell'}}$  implies  $\ell \leq \ell'$ . The corresponding intersection problem will be solved by dealing successively with these summands. The reader will have no difficulty writing out the sets  $I, J, K \subset \{1, 2, \dots, n\}$ , where  $n = r + 2p$ . The following figure illustrates the case  $p = 2$  with  $r = 8$ ,  $a_1 = d_1 = 1$ ,  $b_1 = c_1 = 3$ ,  $a_2 = 2$ ,  $b_2 = c_2 = 1$ , and  $d_2 = 4$ .



We deal next with a somewhat more complicated extremal measure, whose support has the shape pictured below along with its dual.



Denote the attachment projections on the  $A$  side by  $X_1 \leq X_2$ , on the  $B$  side by  $Y_1 \leq Y_2$ , and on the  $C$  side  $Z_1 \leq Z_2$ . In the illustration we used the measure  $m$  which assigns unit mass to the root edges of the skeleton, and this measure has weight  $\omega(m) = 4$ . The measure  $m^*$  is a sum of six extremal measures with supports pictured below.



Denote the corresponding measures in  $\mathcal{M}_4^*$  by  $\mu_\ell$  with  $\ell = 1, 2, \dots, 6$ . the attachment projections for these measures are easily found. For instance,  $\mu_1$  has attachment projections  $X_2^\perp, Y_1^\perp$ , and  $\mu_3$  has attachment projections  $X_2^\perp, Y_2^\perp$ , and  $Z_2^\perp$ . Recalling that clocks run backwards in  $\mathcal{M}^*$ , we easily determine that

$$\mu_1 \prec_0 \mu_4 \prec_0 \mu_6, \quad \mu_2 \prec_0 \mu_5 \prec_0 \mu_6, \quad \mu_3 \prec_0 \mu_6,$$

and no other direct comparisons occur. It is now easy to see that the generic solution is obtained as follows. Form first the projection

$$P_1 = (X_2 \wedge Y_1) \vee (X_2 \wedge Z_2) \vee (X_2 \wedge Y_2 \wedge Z_1)$$


corresponding with the measure  $\mu_1 + \mu_2 + \mu_3$ . Next calculate

$$P_2 = [(X_1 \vee P_1) \wedge (Z_1 \wedge P_1)] \vee [(X_1 \vee P_1) \wedge (Y_2 \wedge P_1)]$$

corresponding with  $\mu_4 + \mu_5$ . Finally, the solution is

$$P = (Y_2 \vee P_2) \wedge (Z_2 \wedge P_2).$$

The examples above illustrate the fact that passing from an extremal measure to its dual yields a dramatic simplification of the intersection problem. We offer, mostly to further illustrate this point, an example of a rather complicated skeleton. The reader will easily identify 15 skeletons in the dual picture.



The number 15 is, not coincidentally, the number of attachment points of the skeleton.

## 7. CONNECTION TO INVARIANT SUBSPACES

The smallest example of sets  $I, J, K$  with  $c_{IJK} > 1$  is given by  $I = J = K = \{2, 4, 6\} \subset \{1, 2, 3, 4, 5, 6\}$ . Fix a  $\text{II}_1$  factor  $\mathcal{A}$  with trace normalized so that  $\tau(1) = 2$ , and fix an element  $T \in \mathcal{A}$ . We will view  $\mathcal{A}$  as an algebra of operators on a Hilbert space  $H$ , and consider the factor  $\mathcal{B} = \mathcal{A} \otimes M_3(\mathbb{C})$  acting on  $H \oplus H \oplus H$ . That is,  $B$  consists of all operator matrices  $[T_{ij}]_{i,j=1}^3$  with  $T_{ij} \in \mathcal{A}$ , with trace defined by

$$\tau([T_{ij}]_{i,j=1}^3) = \sum_{j=1}^3 \tau(T_{jj}).$$

We construct now the following spaces:

$$\begin{aligned} X_2 &= \{\xi \oplus 0 \oplus 0 : \xi \in H\}, \\ Y_2 &= \{0 \oplus \xi \oplus 0 : \xi \in H\}, \\ Z_2 &= \{0 \oplus 0 \oplus \xi : \xi \in H\}, \\ X_4 &= \{\xi \oplus \eta \oplus \eta : \xi, \eta \in H\}, \\ Y_4 &= \{\eta \oplus \xi \oplus \eta : \xi, \eta \in H\}, \\ Z_4 &= \{\eta \oplus T\eta \oplus \xi : \xi, \eta \in H\}. \end{aligned}$$

It is easy to see that the orthogonal projections  $E_2, F_2, G_2, E_4, F_4, G_4$  onto these spaces belong to  $\mathcal{A}$  and

$$\tau(E_j) = \tau(F_j) = \tau(G_j) = j, \quad j = 2, 4.$$

Indeed, we can write these projections explicitly:

$$\begin{aligned} E_2 &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, F_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, G_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \\ E_4 &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix}, F_4 = \begin{bmatrix} \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \end{bmatrix}, \\ G_4 &= \begin{bmatrix} (1 + T^*T)^{-1} & (1 + T^*T)^{-1}T^* & 0 \\ T(1 + T^*T)^{-1} & T(1 + T^*T)^{-1}T^* & 0 \\ 0 & 0 & 1 \end{bmatrix}. \end{aligned}$$

The trace of  $G_4$  is seen to be 4 because  $\begin{bmatrix} (1+T^*T)^{-1} & (1+T^*T)^{-1}T^* \\ T(1+T^*T)^{-1} & T(1+T^*T)^{-1}T^* \end{bmatrix}$  is the range projection of the partial isometry  $\begin{bmatrix} (1+T^*T)^{-1/2} & 0 \\ T(1+T^*T)^{-1/2} & 0 \end{bmatrix}$  which has initial projection  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ . Assume that  $P \in S(\mathcal{E}, I) \cap S(\mathcal{F}, J) \cap S(\mathcal{G}, K)$ . In other words,  $\tau(P) = 3$ ,  $\tau(P \wedge E_2) \geq 1$ ,  $\tau(P \wedge F_2) \geq 1$ ,  $\tau(P \wedge G_2) \geq 1$ ,  $\tau(P \wedge E_4) \geq 2$ ,  $\tau(P \wedge F_4) \geq 2$ , and  $\tau(P \wedge G_4) \geq 2$ . It follows then that there exist projections  $Q, Q', Q'' \in \mathcal{A}$  such that  $\tau(Q) \geq 1$ ,  $\tau(Q') \geq 1$ ,  $\tau(Q'') \geq 1$ , and  $P \geq Q \oplus Q' \oplus Q''$ , which implies that  $\tau(Q) = \tau(Q') = \tau(Q'') = 1$  and  $P = Q \oplus Q' \oplus Q''$ . Next observe that

$$P \wedge E_4 = Q \oplus \left[ (Q' \oplus Q'') \wedge \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix} \right] = Q \oplus \begin{bmatrix} \frac{1}{2}Q' \wedge Q'' & \frac{1}{2}Q' \wedge Q'' \\ \frac{1}{2}Q' \wedge Q'' & \frac{1}{2}Q' \wedge Q'' \end{bmatrix}.$$

This projection must have trace at least 2, and therefore  $Q' = Q''$ . Analogously, the condition  $\tau(P \wedge F_4) \geq 2$  implies that  $Q = Q''$ . We conclude that  $P = Q \oplus Q \oplus Q$ . Finally,  $\tau(P \wedge G_4) \geq 2$  will imply that  $QTQ = TQ$ , so that  $Q$  is an invariant projection for the operator  $T$ . Thus the solution of this particular intersection problem implies the existence of invariant projections of trace 1 for every  $T \in \mathcal{A}$ . In [9] it is shown that this problem has an approximate solution. More precisely, given  $\varepsilon > 0$ , there exist projections  $Q, Q_1 \in \mathcal{A}$  such that  $\tau(Q) = 1$ ,  $Q \leq Q_1$ ,  $\tau(Q_1) < 1 + \varepsilon$ , and  $Q_1 T Q = T Q$ . This leads to an approximate solution of the intersection problem. One would expect that solving the intersection problem for more complicated sets with  $c_{IJK} > 1$  would require considerable progress in the study of  $\text{II}_1$  factors.

## 8. APPLICATIONS OF FREE PROBABILITY

In this brief section we give two applications of free products of von Neumann algebras and free probability. First, we show that all finite von Neumann algebras with a normal, faithful trace admit a trace-preserving embedding into a factor of type  $\text{II}_1$ . This completes the proof of the Horn inequalities for selfadjoint elements in such algebras.

**Proposition 8.1.** *Let  $\mathcal{A}_j$  be von Neumann algebras equipped with normal, faithful, tracial states  $\tau_j$ ,  $j = 1, 2$ , and let  $(\mathcal{A}, \tau) = (\mathcal{A}_1, \tau_1) * (\mathcal{A}_2, \tau_2)$  be the free product von Neumann algebra. If  $\mathcal{A}_2$  is diffuse, i.e., it has no minimal projections, and  $\mathcal{A}_1$  is not a copy of the complex numbers, then  $\mathcal{A}$  is a  $\text{II}_1$  factor.*

*Proof.* Let  $\mathcal{B}$  be the  $C^*$ -subalgebra of  $\mathcal{A}$  generated by the union of the copies of  $\mathcal{A}_1$  and  $\mathcal{A}_2$  in  $\mathcal{A}$ . Then  $\mathcal{B}$  is isomorphic to the  $C^*$ -algebra reduced free product of  $(\mathcal{A}_1, \tau_1)$  and  $(\mathcal{A}_2, \tau_2)$ , and it is dense in  $\mathcal{A}$  in the strong operator topology. By Proposition 3.2 of [12],  $\mathcal{B}$  has a unique tracial state. It follows that  $\mathcal{A}$  has a unique normal tracial state. As  $\mathcal{A}$  is clearly infinite dimensional, it is a  $\text{II}_1$  factor.  $\square$

Next, we will argue that arbitrary projections in a factor of type  $\text{II}_1$  can be perturbed into general position by letting one of them evolve according to free Brownian motion. This perturbation will take place typically in a larger factor obtained as a free product, with the free Brownian motion in one of the factors.

Let  $\mathcal{A}$  be a  $\text{II}_1$  factor with trace  $\tau$ , and let  $P, Q \in \mathcal{A}$  be two projections. Let  $U_t$  be a free right unitary Brownian motion, free from  $\{P, Q\}$ . Recall that a free right

unitary Brownian motion is the solution of the free stochastic differential equation

$$U_0 = 1, \quad dU_t = iU_t dX_t - \frac{1}{2}U_t t dt,$$

where  $X_t$  is a standard additive free Brownian motion (cf. [7]). For our purposes, the following three properties of a unitary Brownian motions are crucial:

- (1)  $t \mapsto U_t$  is norm-continuous;
- (2) for any  $\varepsilon > 0$ ,  $U_t^* U_{t+\varepsilon}$  is free from  $U_s$  for all  $s < t$ .

For the purposes of the following result, we will say that  $P$  and  $Q$  are in general position if  $\tau(P \wedge Q) = \max\{0, \tau(P) + \tau(Q) - 1\}$ .

**Theorem 8.2.** *The projections  $U_t P U_t^*$  and  $Q$  are in general position for every  $t > 0$ .*

*Proof.* Fix  $t > 0$ , and set  $P_t = U_t P U_t^*$ . As in the proof of Proposition 5.3, we may and shall assume that  $\tau(P) + \tau(Q) \leq 1$ . Arguing by contradiction, assume that  $P_t \wedge Q \neq 0$ . Setting  $R = P_t \wedge Q$ , observe that the function

$$f(s) = \tau((R P_s R - R)^2), \quad s \geq 0,$$

is nonnegative, and therefore  $f$  has a minimum at  $s = t$ . The fact that  $U_t$  is a free Brownian motion, and Ito calculus, imply that  $f$  is a differentiable function, and

$$f'(t) = \tau(R)(-1 + \tau(P) + \tau(R) - \tau(R)\tau(P)),$$

where we used the fact that  $(R P_t R)^2 = R P_t R = R$ . Now, we have  $0 < \tau(R) \leq \tau(P)$  and  $1 - \tau(P) \geq \tau(R)$ , so that this relation implies

$$f'(t) \leq -\tau(R)^2 \tau(P) < 0.$$

This however is not compatible with  $f(t)$  being a minimum. □

If  $\mathcal{E}$  and  $\mathcal{F}$  are two flags in  $\mathcal{A}$ , the preceding result yields a unitary  $U$ , arbitrarily close to 1, so that the spaces of the flag  $U\mathcal{F}U^*$  are in general position relative to the spaces of  $\mathcal{E}$ . Dealing with three flags would require the use of two Brownian motions, free from each other and from the flags. In order to obtain flags which are generic for a given intersection of three Schubert cells, this construction would have to be iterated following the inductive procedure of Proposition 5.1.

#### REFERENCES

- [1] P. Belkale, Local systems on  $P^1 \backslash S$  for  $S$  a finite set, *Compositio Math.* **129**(2001), no. 1, 67–86.
- [2] ———, Geometric proofs of Horn and saturation conjectures. *J. Algebraic Geom.* **15**(2006), no. 1, 133–173.
- [3] H. Bercovici and W. S. Li, Inequalities for eigenvalues of sums in a von Neumann algebra. *Recent advances in operator theory and related topics* (Szeged, 1999), 113–126, *Oper. Theory Adv. Appl.*, **127**, Birkhäuser, Basel, 2001.
- [4] ———, Eigenvalue inequalities in an embeddable factor, *Proc. Amer. Math. Soc.* **134**(2006), no. 1, 75–80.
- [5] H. Bercovici, W. S. Li, and T. Smotzer, Continuous versions of the Littlewood-Richardson rule, selfadjoint operators, and invariant subspaces, *J. Operator Theory* **54**(2005), no. 1, 69–92.
- [6] H. Bercovici, W. S. Li, and D. Timotin, The Horn conjecture for sums of compact selfadjoint operators, arXiv:math.FA/0709.1088.
- [7] Ph. Biane, Free Brownian motion, free stochastic calculus and random matrices, *Free probability theory* (Waterloo, ON, 1995), 1–19, Fields Inst. Commun., 12, Amer. Math. Soc., Providence, RI, 1997.



- [8] A. S. Buch, The saturation conjecture (after A. Knutson and T. Tao). With an appendix by William Fulton. *Enseign. Math. (2)* **46**(2000), no. 1-2, 43–60.
- [9] B. Collins and K. Dykema, On a reduction procedure for Horn inequalities in finite von Neumann algebras, arXiv:math.OA/0711.3930.
- [10] ———, A Linearization of Connes' Embedding Problem, arXiv:math.OA/0706.3918
- [11] J. Day, W. So, and R. C. Thompson, The spectrum of a Hermitian matrix sum, *Linear Algebra Appl.* **28** (1998), no. 2-3, 289–332.
- [12] K. Dykema, Simplicity and the stable rank of some free product  $C^*$ -algebras, *Trans. Amer. Math. Soc.* **351**(1999), 1-40.
- [13] S. Friedland, Finite and infinite dimensional generalizations of Klyachko's theorem, *Linear Algebra Appl.* **319**(2000), no. 1-3, 3–22.
- [14] W. Fulton, Eigenvalues of sums of Hermitian matrices (after A. Klyachko), *Séminaire Bourbaki* vol. 1997/1998, Exp. No. 845, *Asterisque* **252**(1998), pp. 255–269.
- [15] ———, Eigenvalues, invariant factors, highest weights, and Schubert calculus, *Bull. Amer. Math. Soc.* **37**(2000), 209–249.
- [16] ———, Eigenvalues of majorized Hermitian matrices and Littlewood-Richardson coefficients, *Linear Algebra Appl.* **319**(2000), 23–36.
- [17] U. Haagerup and H. Schultz, Hanne, Brown measures of unbounded operators affiliated with a finite von Neumann algebra, *Math. Scand.* **100**(2007), no. 2, 209–263.
- [18] ———, Invariant Subspaces for Operators in a General  $II_1$ -factor, arXiv:math.OA/0611256.
- [19] A. Horn, Eigenvalues of sums of Hermitian matrices, *Pacific J. Math.*, **12**(1962), 225–241.
- [20] A. A. Klyachko, Stable bundles, representation theory and Hermitian operators. *Selecta Math. (N.S.)* **4**(1998), no. 3, 419–445.
- [21] A. Knutson and T. Tao, The honeycomb model of  $GL_n(\mathbb{C})$  tensor products. I. Proof of the saturation conjecture, *J. Amer. Math. Soc.* **12**(1999), no. 4, 1055–1090.
- [22] A. Knutson, T. Tao, and C. Woodward, The honeycomb model of  $GL_n(\mathbb{C})$  tensor products. II. Puzzles determine facets of the Littlewood-Richardson cone, *J. Amer. Math. Soc.* **17**(2004), no. 1, 19–48.
- [23] R. C. Thompson and L. J. Freede, On the eigenvalues of sums of Hermitian matrices, *Linear Algebra and Appl.* **4**(1971), 369–376.
- [24] ———, On the eigenvalues of sums of Hermitian matrices. II, *Aequationes Math.* **5**(1970), 103–115.
- [25] R. C. Thompson and S. Therianos, On a construction of B. P. Zwahlen. *Linear and Multilinear Algebra* **1**(1973/74), 309–325.
- [26] R. Vakil, A geometric Littlewood-Richardson rule. Appendix A written with A. Knutson, *Ann. of Math. (2)* **164**(2006), no. 2, 371–421.
- [27] ———, Schubert induction, *Ann. of Math. (2)* **164**(2006), no. 2, 489–512.

HB: DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405, USA  
*E-mail address:* `bercovic@indiana.edu`

BC: DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, OTTAWA, ONTARIO K1N 6N5 CANADA, AND CNRS, DEPARTMENT OF MATHEMATICS, UNIVERSITÉ CLAUDE BERNARD, LYON 1, LYON, FRANCE  
*E-mail address:* `bcollins@uottawa.edu`

KD: DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA  
*E-mail address:* `kdykema@math.tamu.edu`

WSL: SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-1060, USA  
*E-mail address:* `li@math.gatech.edu`

DT: SIMION STOILOW INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, PO Box 1-764, BUCHAREST 014700, ROMANIA  
*E-mail address:* `Dan.Timotin@imar.ro`